

Example 2.

Find the solution of (4) that satisfies the auxiliary condition $u(0, y) = y^3$. Indeed, putting $x = 0$ in (7), we get $y^3 = f(e^{-0}y)$, so that $f(y) = y^3$. Therefore, $u(x, y) = (e^{-x}y)^3 = e^{-3x}y^3$. \square

Example 3.

Solve the PDE

$$u_x + 2xy^2u_y = 0. \quad (8)$$

The characteristic curves satisfy the ODE $dy/dx = 2xy^2/1 = 2xy^2$. To solve the ODE, we separate variables: $dy/y^2 = 2x dx$; hence $-1/y = x^2 - C$, so that

$$y = (C - x^2)^{-1}. \quad (9)$$

These curves are the characteristics. Again, $u(x, y)$ is a constant on each such curve. (Check it by writing it out.) So $u(x, y) = f(C)$, where f is an arbitrary function. Therefore, the general solution of (8) is obtained by solving (9) for C . That is,

$$u(x, y) = f\left(x^2 + \frac{1}{y}\right). \quad (10)$$

Again this is easily checked by differentiation, using the chain rule: $u_x = 2x \cdot f'(x^2 + 1/y)$ and $u_y = -(1/y^2) \cdot f'(x^2 + 1/y)$, whence $u_x + 2xy^2u_y = 0$. \square

In summary, the geometric method works nicely for any PDE of the form $a(x, y)u_x + b(x, y)u_y = 0$. It reduces the solution of the PDE to the solution of the ODE $dy/dx = b(x, y)/a(x, y)$. If the ODE can be solved, so can the PDE. Every solution of the PDE is constant on the solution curves of the ODE.

Moral Solutions of PDEs generally depend on arbitrary functions (instead of arbitrary constants). You need an auxiliary condition if you want to determine a unique solution. Such conditions are usually called *initial* or *boundary* conditions. We shall encounter these conditions throughout the book.

EXERCISES

1. Solve the first-order equation $2u_t + 3u_x = 0$ with the auxiliary condition $u = \sin x$ when $t = 0$.
2. Solve the equation $3u_y + u_{xy} = 0$. (*Hint*: Let $v = u_y$.)

10 CHAPTER 1 WHERE PDEs COME FROM

3. Solve the equation $(1 + x^2)u_x + u_y = 0$. Sketch some of the characteristic curves.
4. Check that (7) indeed solves (4).
5. Solve the equation $xu_x + yu_y = 0$.
6. Solve the equation $\sqrt{1 - x^2}u_x + u_y = 0$ with the condition $u(0, y) = y$.
7. (a) Solve the equation $yu_x + xu_y = 0$ with $u(0, y) = e^{-y^2}$.
(b) In which region of the xy plane is the solution uniquely determined?
8. Solve $au_x + bu_y + cu = 0$.
9. Solve the equation $u_x + u_y = 1$.
10. Solve $u_x + u_y + u = e^{x+2y}$ with $u(x, 0) = 0$.
11. Solve $au_x + bu_y = f(x, y)$, where $f(x, y)$ is a given function. If $a \neq 0$, write the solution in the form

$$u(x, y) = (a^2 + b^2)^{-1/2} \int_L f ds + g(bx - ay),$$

where g is an arbitrary function of one variable, L is the characteristic line segment from the y axis to the point (x, y) , and the integral is a line integral. (*Hint*: Use the coordinate method.)

12. Show that the new coordinate axes defined by (3) are orthogonal.
13. Use the coordinate method to solve the equation

$$u_x + 2u_y + (2x - y)u = 2x^2 + 3xy - 2y^2.$$

1.3 FLOWS, VIBRATIONS, AND DIFFUSIONS

The subject of PDEs was practically a branch of physics until the twentieth century. In this section we present a series of examples of PDEs as they occur in physics. They provide the basic motivation for all the PDE problems we study in the rest of the book. We shall see that most often in physical problems the independent variables are those of space x, y, z , and time t .

Example 1. Simple Transport

Consider a fluid, water, say, flowing at a constant rate c along a horizontal pipe of fixed cross section in the positive x direction. A substance, say a pollutant, is suspended in the water. Let $u(x, t)$ be its concentration in grams/centimeter at time t . Then

$$u_t + cu_x = 0. \tag{1}$$

(That is, the rate of change u_t of concentration is proportional to the gradient u_x . Diffusion is assumed to be negligible.) Solving this equation as in Section 1.2, we find that the concentration is a function of $(x - ct)$