

# Notes on Differential Geometry

Fernando Manfio  
University of São Paulo

2024

# Contents

<b>1</b>	<b>Regular surfaces</b>	<b>1</b>
1.1	Regular surfaces . . . . .	1
1.2	Differentiable mappings between surfaces . . . . .	5
1.3	The tangent plane . . . . .	8
1.4	Orientable surfaces . . . . .	11
<b>2</b>	<b>The local theory of surfaces</b>	<b>17</b>
2.1	Derivatives of vector fields in $\mathbb{R}^3$ . . . . .	17
2.2	Derivatives of vector fields on surfaces . . . . .	22
2.3	The Gauss map of a regular surface . . . . .	26
2.4	Euler's principal curvatures . . . . .	29
2.5	The curvature of a regular surface . . . . .	31
2.6	The curvature in local coordinates . . . . .	33
2.7	Rotational surfaces . . . . .	37
2.8	Minimal surfaces . . . . .	39
2.9	Normal variations . . . . .	43
2.10	Appendix: Regular curves . . . . .	47
	<b>Bibliography</b>	<b>50</b>

# Chapter 1

## Regular surfaces

In these notes the term differentiable means differentiable of class  $C^\infty$ .

### 1.1 Regular surfaces

In few words, a regular surface in  $\mathbb{R}^3$  is a subset that is locally homeomorphic to an open subset of  $\mathbb{R}^2$ .

**Definition 1.1.1.** A subset  $M \subset \mathbb{R}^3$  is called a *regular surface* if, for each point  $p \in M$ , there exists an open set  $V \subset \mathbb{R}^3$ , with  $p \in V$ , and a homeomorphism  $\varphi : U \rightarrow M \cap V$  defined on an open set  $U \subset \mathbb{R}^2$ , such that

- (a)  $\varphi$  is differentiable,
- (b) For each point  $x \in U$ , the differential  $d\varphi(x) : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  is injective.

The mapping  $\varphi$  is called a *parametrization* of  $M$  around  $p$ , and the subset  $M \cap V$  is called a *coordinate neighborhood* of  $M$ . This means that  $M$  is endowed with the induced topology of  $\mathbb{R}^3$ , and therefore any regular surface is, in particular, a topological subspace of  $\mathbb{R}^3$ .

The condition that  $\varphi$  is differentiable means that if we write

$$\varphi(x_1, x_2) = (\varphi_1(x_1, x_2), \varphi_2(x_1, x_2), \varphi_3(x_1, x_2)),$$

then the coordinate functions  $\varphi_1, \varphi_2, \varphi_3$  have continuous partial derivatives of all orders in the open set  $U$ .

The condition that  $d\varphi(x) : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  is a injective linear map is equivalent to any of the following conditions:

- (a) The set  $\{d\varphi(x) \cdot e_i : 1 \leq i \leq 2\}$  is linearly independent, where  $\{e_1, e_2\}$  denotes the canonical basis of  $\mathbb{R}^2$ .
- (b) The Jacobian matrix  $d\varphi(x)$  has rank two at any point  $x \in U$ .

**Example 1.1.2.** Any two-dimensional vector space  $E \subset \mathbb{R}^3$  is a regular surface. In fact, consider a linear isomorphism  $T: E \rightarrow \mathbb{R}^2$ , and endow  $E$  with the unique topology (induced of  $\mathbb{R}^3$ ) that makes  $T$  a homeomorphism. Since any linear mapping into  $\mathbb{R}^2$  is differentiable, it follows that  $T$  is a diffeomorphism, and thus  $T$  is a global parametrization of  $E$ .

**Example 1.1.3.** Let us show that the unit sphere  $\mathbb{S}^2 \subset \mathbb{R}^3$  is a regular surface. Fix a point  $p \in \mathbb{S}^2$  other than the north pole  $N = (0, 0, 1)$  and consider the stereographic projection  $\pi_N: \mathbb{S}^2 \setminus \{N\} \rightarrow \mathbb{R}^2$ . We already know that  $\pi_N$  is a homeomorphism, whose inverse is the map  $\varphi: \mathbb{R}^2 \rightarrow \mathbb{S}^2 \setminus \{N\}$  given by

$$\varphi(x) = \left( \frac{2x_1}{\|x\|^2 + 1}, \frac{2x_2}{\|x\|^2 + 1}, \frac{\|x\|^2 - 1}{\|x\|^2 + 1} \right),$$

for every point  $x = (x_1, x_2) \in \mathbb{R}^2$ . Since each coordinate function of  $\varphi$  is differentiable, it follows that  $\varphi$  is also differentiable. It is straightforward to check that  $d\varphi(x)$  has rank two at any point  $x \in \mathbb{R}^2$ . Finally, if  $p = N$ , just consider the stereographic projection  $\pi_S$  relative to the south pole  $S \in \mathbb{S}^2$ .

**Example 1.1.4.** The graph of a differentiable function  $f: U \rightarrow \mathbb{R}$ , defined in an open set  $U \subset \mathbb{R}^2$ , is a regular surface. In fact, denoting by  $\text{Gr}(f)$  the graph of  $f$ , let us show that the map  $\varphi: U \rightarrow \mathbb{R}^3$  given by

$$\varphi(x) = (x, f(x)),$$

is a global parametrization of  $\text{Gr}(f)$ . Since  $f$  is differentiable, the same holds for  $\varphi$ . Each point  $(x, f(x)) \in \text{Gr}(f)$  is the image under  $\varphi$  of the unique point  $x \in U$ , and  $\varphi$  is therefore injective. Moreover, the restriction to  $\text{Gr}(f)$  of the projection of  $\mathbb{R}^3$  onto  $\mathbb{R}^2$  is a inverse to  $\varphi$ , and this shows that  $\varphi^{-1}$  is also continuous. It follows that  $\varphi$  is a homeomorphism. Finally, it is easy to see that  $d\varphi(x)$  has rank two at any point  $x \in U$ .

The following result provides a local converse of Example 1.1.4. More precisely, any regular surface is locally the graph of a differentiable function.

**Proposition 1.1.5.** Given a regular surface  $M \subset \mathbb{R}^3$  and a point  $p \in M$ , there exist an open set  $U \subset \mathbb{R}^2$ , an open set  $V \subset \mathbb{R}^3$  with  $p \in V$ , and a differentiable function  $g: U \rightarrow \mathbb{R}$  such that  $M \cap V = \text{Gr}(g)$ .

*Proof.* Fix a point  $p \in M$  and consider a parametrization  $\varphi : U \rightarrow \varphi(U)$  of  $M$ , with  $p = \varphi(x)$ . Since  $E = d\varphi(x)(\mathbb{R}^2)$  is a two-dimensional vector subspace of  $\mathbb{R}^3$ , there exists an orthogonal decomposition  $\mathbb{R}^3 = \mathbb{R}^2 \oplus \mathbb{R}$  such that the projection  $\pi : \mathbb{R}^2 \times \mathbb{R} \rightarrow \mathbb{R}^2$  maps  $E$  isomorphically onto  $\mathbb{R}^2$ , and define the map

$$\eta = \pi \circ \varphi : U \rightarrow \mathbb{R}^2.$$

Since  $d\eta(q) = \pi \circ d\varphi(q)$  is a linear isomorphism, it follows from the inverse function theorem that there exist a open set  $W \subset \mathbb{R}^2$ , with  $q \in W \subset U$ , such that  $\eta|_W : W \rightarrow \eta(W) = Z$  is a diffeomorphism. Define

$$\xi = (\eta|_W)^{-1} : Z \rightarrow W \quad \text{and} \quad \psi = \varphi \circ \xi.$$

It follows that  $\psi$  is also a parametrization of  $M$  and

$$\pi \circ \psi = \pi \circ (\varphi \circ \xi) = \eta \circ \xi = id.$$

According to the orthogonal decomposition  $\mathbb{R}^3 = \mathbb{R}^2 \oplus \mathbb{R}$ , it follows from the above equality that the first coordinate of  $\psi(x)$  is  $x$ . Let us denote by  $g(x)$  the second one. Thus,

$$\psi(Z) = \varphi(W) = \{(x, g(x)) : x \in W\}$$

for some differentiable function  $g : W \rightarrow \mathbb{R}$ . Since  $\varphi$  is a open map, one has

$$\varphi(W) = M \cap V = \text{Gr}(g),$$

for some open set  $V \subset \mathbb{R}^3$ , with  $p \in V$ . □

Let us look a simple application of Proposition 1.1.5.

**Example 1.1.6.** Let us consider the *one-sheeted cone*  $M \subset \mathbb{R}^3$  given by

$$M = \{(x, y, z) : x^2 + y^2 = z^2, z \geq 0\}.$$

We will show that  $M$  is not a regular surface. If  $M$  were a regular surface then, by virtue of Proposition 1.1.5,  $M$  would be locally a graph of a differential function around  $(0, 0, 0)$ . More precisely, there exist open sets  $U \subset \mathbb{R}^2$  and  $V \subset \mathbb{R}^3$ , with  $0 \in V$ , and a differentiable function  $g : U \rightarrow \mathbb{R}$  such that  $M \cap V = \text{Gr}(g)$ . Note that, according to a decomposition  $\mathbb{R}^3 = \mathbb{R}^2 \oplus \mathbb{R}$ , the only possibility for  $M \cap V$  to be a graph is for the second factor to be the axis- $z$ . Thus, it follows that  $g = f|_U$ , where  $f(x, y) = \sqrt{x^2 + y^2}$ . However,  $f$  is not differentiable at  $(0, 0)$ .

Let  $f : V \rightarrow \mathbb{R}$  be a differentiable function, defined in an open set  $V \subset \mathbb{R}^3$ . We say that a point  $p \in V$  is a *regular point* of  $f$  if the differential  $df(p)$  is surjective, that is,  $df(p) \neq 0$ . A point  $c \in \mathbb{R}$  is called a *regular value* of  $f$  if the inverse image  $f^{-1}(c)$  contains only regular points of  $f$ . Notice that any point  $c \notin f(V)$  is trivially a regular value of  $f$ .

**Proposition 1.1.7.** Let  $f : V \rightarrow \mathbb{R}$  be a differentiable function, defined in an open set  $V \subset \mathbb{R}^3$ , and  $c \in \mathbb{R}$  be a regular value of  $f$ . If  $f^{-1}(c) \neq \emptyset$ , then  $M = f^{-1}(c)$  is a regular surface.

*Proof.* By virtue of Example 1.1.4, it suffices to prove that  $M$  is locally graph of some differentiable function. Given a point  $p \in M$ , with  $p = (x_0, y_0, z_0)$ , we can assume that  $\frac{\partial f}{\partial z}(p) \neq 0$ . Therefore, it follows from the implicit function theorem that there exist an open set  $W = U \times I$ , where  $U$  is an open set of  $\mathbb{R}^2$  with  $(x_0, y_0) \in U$ , and  $I$  is an open interval with  $z_0 \in I$ , and a differentiable function  $g : U \rightarrow \mathbb{R}$  such that

$$f((x, y), g(x, y)) = c,$$

for every  $(x, y) \in U$ . This proves that  $M \cap W = \text{Gr}(g)$ .  $\square$

**Example 1.1.8.** The unit sphere  $\mathbb{S}^2 \subset \mathbb{R}^3$  can be described as the inverse image  $f^{-1}(1)$  of the function  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$  given by

$$f(x) = \|x\|^2 = \langle x, x \rangle,$$

for every  $x \in \mathbb{R}^3$ . Notice that  $f$  is differentiable and, for any point  $p \in \mathbb{R}^3$  and any vector  $v \in \mathbb{R}^3$ , we obtain

$$df(p) \cdot v = 2\langle p, v \rangle.$$

This implies that  $0 \in \mathbb{R}^3$  is the unique critical point of  $f$ . Since  $f(0) = 0 \neq 1$ , we conclude that 1 is regular value of  $f$ . Therefore, the sphere  $\mathbb{S}^2$  is a regular surface as we have already seen.

**Remark 1.1.9.** The inverse image  $f^{-1}(c)$  can be a regular surface without  $c$  being a regular value of  $f$ . For instance, consider the function  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$  given by  $f(x, y, z) = z^2$ . Note that  $f$  is differentiable and  $f^{-1}(0)$  is the plane- $xy$ , which is a regular surface in  $\mathbb{R}^3$ . However, the point  $0 \in \mathbb{R}$  is not a regular value of  $f$ , because  $df(x, y, 0) = 0$ , for every  $(x, y, 0) \in f^{-1}(0)$ .

The following result is a converse of Proposition 1.1.7.

**Theorem 1.1.10.** *Any regular surface  $M$  in  $\mathbb{R}^3$  is locally the inverse image of a regular value. More precisely, given a point  $p \in M$ , there exist an open set  $V \subset \mathbb{R}^3$  with  $p \in V$ , and a differentiable function  $f: V \rightarrow \mathbb{R}$  such that  $M \cap V = f^{-1}(0)$ , where  $0 \in \mathbb{R}$  is a regular value of  $f$ .*

*Proof.* It follows from Proposition 1.1.5 that there exists an open set  $V \subset \mathbb{R}^3$ , with  $p \in V$ , such that  $M \cap V = \text{Gr}(g)$ , where  $g: U \rightarrow \mathbb{R}$  is a differentiable function defined in an open set  $U \subset \mathbb{R}^2$ . Define a function  $f: V \rightarrow \mathbb{R}$  by  $f(x, y) = y - g(x)$ . By construction, one has

$$M \cap V = \text{Gr}(g) = f^{-1}(0).$$

It suffices to prove that  $df(x, y)$  is surjective at any point  $(x, y) \in f^{-1}(0)$ . In fact, given  $(x, y) \in f^{-1}(0)$  and  $(u, v) \in \mathbb{R}^3$ , we obtain:

$$\begin{aligned} df(x, y) \cdot (u, v) &= df(x, y) \cdot (u, 0) + df(x, y) \cdot (0, v) \\ &= \text{Id}(0) - dg(x) \cdot u + \text{Id}(v) - dg(x) \cdot 0 \\ &= v - dg(x) \cdot u. \end{aligned}$$

Therefore, given  $v \in \mathbb{R}$ , one has  $df(x, y) \cdot (0, v) = v$ , and this proves that  $0$  is a regular value of  $f$ .  $\square$

## 1.2 Differentiable mappings between surfaces

In this section we will define what it means for a map  $f: M \rightarrow N$ , between two regular surfaces  $M$  and  $N$ , to be differentiable at a point  $p \in M$ .

**Definition 1.2.1.** A map  $f: M \rightarrow N$ , between the regular surfaces  $M$  and  $N$ , is said to be *differentiable* at a point  $p \in M$  if there exist parametrizations  $\varphi: U \rightarrow \varphi(U)$  of  $M$  and  $\psi: V \rightarrow \psi(V)$  of  $N$ , with  $f(\varphi(U)) \subset \psi(V)$  and  $p = \varphi(x)$ , such that the map

$$\psi^{-1} \circ f \circ \varphi: U \rightarrow V \tag{1.1}$$

is differentiable at  $x \in U$ .

The map given in (1.1) is called a *representation* of  $f$  in terms of the parametrizations  $\varphi$  and  $\psi$ . We have to show that this definition does not depend on the choice of parametrizations. In fact, consider parametrizations  $\varphi': U' \rightarrow \varphi'(U')$  of  $M$  and  $\psi': V' \rightarrow \psi'(V')$  of  $N$ , with  $p \in \varphi'(U')$  and  $f(\varphi'(U')) \subset \psi'(V')$ . In the intersection  $\varphi'^{-1}(\varphi(U) \cap \varphi'(U'))$ , one has

$$\psi'^{-1} \circ f \circ \varphi' = (\psi'^{-1} \circ \psi) \circ (\psi^{-1} \circ f \circ \varphi) \circ (\varphi^{-1} \circ \varphi').$$

Therefore, the differentiability of  $f$  will be well-defined if we prove that  $\psi'^{-1} \circ \psi$  and  $\varphi^{-1} \circ \varphi'$  are differentiable. But this will be a consequence of the following lemma.

Let  $\varphi : U \rightarrow \varphi(U)$  and  $\psi : V \rightarrow \psi(V)$  be two parametrizations of a regular surface  $M$  such that  $\varphi(U) \cap \psi(V) \neq \emptyset$ . The map

$$\psi^{-1} \circ \varphi : \varphi^{-1}(W) \rightarrow \psi^{-1}(W), \quad (1.2)$$

is called the *change of coordinates* between the parametrizations  $\varphi$  and  $\psi$ , where  $W = \varphi(U) \cap \psi(V)$ .

**Lemma 1.2.2.** The change of coordinates (1.2) is a diffeomorphism.

*Proof.* Fix a point  $p \in \varphi(U) \cap \psi(V)$ , with  $p = \varphi(q)$ . Since  $d\varphi(q)$  is injective, and writing

$$\varphi(u, v) = (x(u, v), y(u, v), z(u, v)),$$

we can assume without loss of generality that  $\frac{\partial(x, y)}{\partial(u, v)}(q) \neq 0$ . Define a map  $\xi : U \times \mathbb{R} \rightarrow \mathbb{R}^3$  by

$$\xi(u, v, w) = (x(u, v), y(u, v), z(u, v) + w).$$

$\xi$  is clearly differentiable and  $\xi|_{U \times \{0\}} = \varphi$ , thus

$$\det(d\xi(q, 0)) = \det \begin{pmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & 0 \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & 0 \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & 1 \end{pmatrix} = \frac{\partial(x, y)}{\partial(u, v)}(q) \neq 0.$$

Therefore, it follows from inverse function theorem that there exists an open set  $K \subset \mathbb{R}^3$ , with  $(q, 0) \in K$ , such that  $\xi|_K : K \rightarrow \xi(K)$  is a diffeomorphism. Note that  $Z = \xi(K)$  is an open set of  $\mathbb{R}^3$ , with  $p \in Z$ . Since  $\xi|_{U \times \{0\}} = \varphi$ , one has

$$\xi^{-1}|_{\varphi(U) \cap Z} = \varphi^{-1}|_{\varphi(U) \cap Z}.$$

On the other hand, since  $\varphi(U) \cap Z$  is an open set of  $M$  and  $\psi$  is a homeomorphism, it follows that  $\psi^{-1}(\varphi(U) \cap Z)$  is an open set of  $V$ . Therefore,

$$\varphi^{-1} \circ \psi|_{\psi^{-1}(\varphi(U) \cap Z)} = \xi^{-1} \circ \psi|_{\psi^{-1}(\varphi(U) \cap Z)}$$

is differentiable as a composition of differentiable maps. Analogously, we can show that  $\psi^{-1} \circ \varphi$  is differentiable, and thus it is a diffeomorphism.  $\square$

Let us explore some consequences.

**Corollary 1.2.3.** Let  $M, N$  be regular surfaces, and assume that  $M \subset V$ , where  $V$  is an open set of  $\mathbb{R}^3$ , and that  $f: V \rightarrow \mathbb{R}^3$  is a differentiable map such that  $f(M) \subset N$ . Then the restriction  $f|_M: M \rightarrow N$  is a differentiable map.

*Proof.* Given a point  $p \in M$ , consider parametrizations  $\varphi: U \rightarrow \varphi(U)$  of  $M$  and  $\psi: V \rightarrow \psi(V)$  of  $N$ , with  $p \in \varphi(U)$  and  $f(\varphi(U)) \subset \psi(V)$ . Then, the map

$$\psi^{-1} \circ f \circ \varphi: U \rightarrow V$$

is differentiable. □

**Example 1.2.4.** The map  $f: \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by  $f(x, y, z) = (ax, by, cz)$ , where  $a, b, c$  are positive real numbers, is clearly differentiable. The restriction of  $f$  to the unit sphere  $\mathbb{S}^2 \subset \mathbb{R}^3$  is differentiable. In fact,  $f|_{\mathbb{S}^2}$  is a differentiable map of the sphere  $\mathbb{S}^2$  into the ellipsoid  $\mathcal{E}$ .

**Remark 1.2.5.** In the case of a map  $f: M \rightarrow \mathbb{R}^2$ , of a regular surface  $M$  into  $\mathbb{R}^2$ , the Definition 1.2.1 takes a rather simpler form. Namely, in this case,  $f$  is differentiable at  $p \in M$  if there exists a parametrization  $\varphi: U \rightarrow \varphi(U)$  of  $M$ , with  $p = \varphi(x)$ , such that

$$f \circ \varphi: U \rightarrow \mathbb{R}^2$$

is differentiable at  $x \in U$ . In fact, just consider  $\psi$  equal to the identity in Definition 1.2.1.

**Corollary 1.2.6.** If  $\varphi: U \rightarrow \varphi(U)$  is a parametrization of a regular surface  $M$ , then  $\varphi^{-1}: \varphi(U) \rightarrow \mathbb{R}^2$  is also differentiable.

*Proof.* Given a point  $p \in \varphi(U)$ , consider the parametrization  $\varphi: U \rightarrow \varphi(U)$  of  $M$ . One has  $p \in \varphi(U)$  and the representation of  $\varphi^{-1}$  in terms of  $\varphi$  is just the identity map, which is differentiable. □

Two regular surfaces  $M$  and  $N$  are *diffeomorphic* if there exists a bijective differentiable map  $f: M \rightarrow N$ , whose inverse  $f^{-1}: N \rightarrow M$  is also differentiable. In this case,  $f$  is called a *diffeomorphism* from  $M$  to  $N$ . In particular, it follows from Corollary 1.2.6 that, if  $\varphi: U \rightarrow \varphi(U)$  is a parametrization of a regular surface  $M$ , then  $U$  and  $\varphi(U)$  are diffeomorphic.

Finally, we will now give a definition for a differentiable function on a regular surface.

**Definition 1.2.7.** A function  $f: M \rightarrow \mathbb{R}$ , defined on a regular surface  $M$ , is said to be *differentiable* at a point  $p \in M$  if there exists a parametrization  $\varphi: U \rightarrow \varphi(U)$  of  $M$ , with  $p = \varphi(x)$ , such that the composition

$$f \circ \varphi: U \rightarrow \mathbb{R}$$

is differentiable at  $x \in U$ .

It follows from Lemma 1.2.2 that the Definition 1.2.7 does not depend on the choice of the parametrization  $\varphi$ . In fact, if  $\psi: V \rightarrow \psi(V)$  is another parametrization of  $M$ , with  $p = \psi(y)$ , then

$$f \circ \psi = (f \circ \varphi) \circ (\varphi^{-1} \circ \psi)$$

is also differentiable.

**Corollary 1.2.8.** If  $f: \mathbb{R}^3 \rightarrow \mathbb{R}$  is a differentiable function, then the restriction of  $f$  to any regular surface  $M$  is a differentiable function on  $M$ .

*Proof.* For any point  $p \in M$  and any parametrization  $\varphi: U \rightarrow \varphi(U)$  of  $M$ , with  $p = \varphi(x)$ , the function  $f \circ \varphi: U \rightarrow \mathbb{R}$  is differentiable at  $x \in U$ .  $\square$

**Example 1.2.9.** Given a regular surface  $M$  and a unit vector  $v \in \mathbb{R}^3$ , consider the height function  $f: M \rightarrow \mathbb{R}$  relative to  $v$ , given by  $f(p) = \langle p, v \rangle$  for every  $p \in M$ . It follows immediately from Corollary 1.2.8 that  $f$  is differentiable.

### 1.3 The tangent plane

In this section we will define the tangent plane to a regular surface  $M$  at a point  $p \in M$ . Before stating the concepts we will need some terminology.

A *parametrized differentiable curve* in  $\mathbb{R}^3$  is just a differentiable map  $\alpha: I \rightarrow \mathbb{R}^3$  defined in an open interval  $I \subset \mathbb{R}$ . The term differentiable means that  $\alpha$  is a correspondence which maps each instant  $t \in I$  into a point  $\alpha(t) = (x(t), y(t), z(t)) \in \mathbb{R}^3$  in such way that the functions  $x(t), y(t), z(t)$  are differentiable. The variable  $t$  is called the *parameter* of the curve. The vector  $\alpha'(t) = (x'(t), y'(t), z'(t)) \in \mathbb{R}^3$  is called the *tangent vector* of  $\alpha$  at  $t$ . The image set  $\alpha(I) \subset \mathbb{R}^3$  is called the *trace* of  $\alpha$ .

Let  $M$  be a regular surface. A differentiable curve  $\alpha: I \rightarrow M$  is simply a differentiable curve  $\alpha: I \rightarrow \mathbb{R}^3$  such that  $\alpha(I) \subset M$ , that is,  $\alpha(t) \in M$  for every  $t \in I$ . Fix a point  $p \in M$ . A vector  $v \in \mathbb{R}^3$  is called a *tangent vector*

to  $M$  at  $p$  if there exists a differentiable curve  $\alpha: (-\epsilon, \epsilon) \rightarrow M$  such that  $\alpha(0) = p$  and  $\alpha'(0) = v$ . The *tangent plane* to  $M$  at  $p$  is the collection of all tangent vectors to  $M$  at  $p$ , and it will be denoted by  $T_p M$ .

**Proposition 1.3.1.** For any parametrization  $\varphi: U \rightarrow \varphi(U)$  of  $M$ , with  $p = \varphi(x)$ , one has

$$T_p M = d\varphi(x)(\mathbb{R}^2).$$

*Proof.* Any vector in the image of  $d\varphi(x)$  is of the form  $d\varphi(x) \cdot v$ , for some  $v \in \mathbb{R}^2$  and therefore is the tangent vector at 0 of the differentiable curve  $\alpha(t) = \varphi(x + tv)$ . Conversely, let  $v \in T_p M$ , with  $v = \alpha'(0)$ , where  $\alpha: (-\epsilon, \epsilon) \rightarrow M$  is a differentiable curve, with  $\alpha(0) = p$ . By virtue of Corollary 1.2.6, the curve  $\gamma = \varphi^{-1} \circ \alpha: (-\epsilon, \epsilon) \rightarrow U$  is differentiable, with  $\beta(0) = x$ . Since  $\alpha = \varphi \circ \beta$ , it follows from the chain rule that

$$v = \alpha'(0) = d\varphi(x) \cdot \gamma'(0)$$

lies in the image of  $d\varphi(x)$ . □

It follows directly from Proposition 1.3.1 that the tangent plane  $T_p M$  is a two-dimensional vector subspace of  $\mathbb{R}^3$ , and it does not depend on the parametrization  $\varphi$ . Moreover, the choice of a parametrization  $\varphi: U \rightarrow \varphi(U)$  of  $M$ , with  $p = \varphi(q)$ , determines a basis  $\{\varphi_u(p), \varphi_v(p)\}$  of  $T_p M$ , called the *basis associated to  $\varphi$* . Here the notations  $\varphi_u, \varphi_v$  mean

$$\varphi_u(p) = \frac{\partial \varphi}{\partial u}(q) = d\varphi(q) \cdot e_1 \quad \text{and} \quad \varphi_v(p) = \frac{\partial \varphi}{\partial v}(q) = d\varphi(q) \cdot e_2$$

Let us see how to determine the coordinates of a vector  $v \in T_p M$  in the basis  $\{\varphi_u(p), \varphi_v(p)\}$  associated to a parametrization  $\varphi: U \rightarrow \varphi(U)$  of  $M$ , with  $p = \varphi(q)$ .

**Example 1.3.2.** Let  $M$  be a regular surface given as the inverse image under a differentiable function  $f: \mathbb{R}^3 \rightarrow \mathbb{R}$  of a regular value, namely  $M = f^{-1}(c)$ . We claim that

$$T_p M = \ker df(p)$$

for any  $p \in M$ . In fact, let  $w \in T_p M$ , with  $w = \alpha'(0)$ , where  $\alpha: (-\epsilon, \epsilon) \rightarrow M$  is a differentiable curve, with  $\alpha(0) = p$ . Then,  $\beta(t) = (f \circ \alpha)(t)$  is a constant curve along  $(-\epsilon, \epsilon)$ . By the chain rule we obtain

$$0 = \beta'(0) = df(p) \cdot w,$$

and this proves the inclusion  $T_p M \subset \ker df(p)$  and hence the equality by dimensional reasons.

Given a differentiable map  $f: M \rightarrow N$  and a point  $p \in M$ , we want to define the differential of  $f$  at  $p$ , denoted by  $df(p)$ , and being a linear map from  $T_p M$  to  $T_{f(p)} N$ . More precisely, for each vector  $w \in T_p M$ , define

$$df(p) \cdot w = (f \circ \alpha)'(0), \quad (1.3)$$

where  $\alpha: (-\epsilon, \epsilon) \rightarrow M$  is a differentiable curve with  $\alpha(0) = p$  and  $\alpha'(0) = w$ .

**Proposition 1.3.3.** The map  $df(p): T_p M \rightarrow T_{f(p)} N$  given in (1.3) is well-defined and is linear, and it will be called the *differential* of  $f$  at  $p \in M$ .

*Proof.* Firstly, we have to check that  $df(p) \cdot w$  does not depend on the choice of curve  $\alpha$ . Let  $\varphi: U \rightarrow \varphi(U)$  and  $\psi: V \rightarrow \psi(V)$  parametrizations of  $M$  and  $N$ , respectively, with  $p = \varphi(q)$  and  $f(\varphi(U)) \subset \psi(V)$ . Writing

$$\varphi = \varphi(u, v) \quad \text{and} \quad \psi = \psi(z, w),$$

suppose that  $f$  is expressed in these coordinates by

$$f(u, v) = (f_1(u, v), f_2(u, v))$$

and that  $\alpha$  is expressed by

$$\alpha(t) = (u(t), v(t)).$$

Thus, the curve  $\beta = f \circ \alpha$  can be write as

$$\beta(t) = (f_1(u(t), v(t)), f_2(u(t), v(t))),$$

and the expression of  $\beta'(0)$  in the basis  $\{\psi_z, \psi_w\}$  is

$$\beta'(0) = \left( \frac{\partial f_1}{\partial u} u'(0) + \frac{\partial f_1}{\partial v} v'(0), \frac{\partial f_2}{\partial u} u'(0) + \frac{\partial f_2}{\partial v} v'(0) \right). \quad (1.4)$$

The relation (1.4) shows that  $\beta'(0)$  depends only on the map  $f$  and the coordinates  $(u'(0), v'(0))$  of  $w$  in the basis  $\{\varphi_u, \varphi_v\}$ . Therefore,  $\beta'(0)$  is independent of  $\alpha$ . Moreover, it also follows from (1.4) that

$$\beta'(0) = df(p) \cdot w = \begin{pmatrix} \frac{\partial f_1}{\partial u} & \frac{\partial f_1}{\partial v} \\ \frac{\partial f_2}{\partial u} & \frac{\partial f_2}{\partial v} \end{pmatrix} \begin{pmatrix} u'(0) \\ v'(0) \end{pmatrix}. \quad (1.5)$$

This shows that  $df(p)$  is a linear map from  $T_p M$  to  $T_{f(p)} N$ , whose matrix in the basis  $\{\varphi_u, \varphi_v\}$  of  $T_p M$  and  $\{\psi_z, \psi_w\}$  of  $T_{f(p)} N$  is just the matrix given in (1.5).  $\square$

Analogously, given a differentiable function  $f: M \rightarrow \mathbb{R}$ , we can define the differential of  $f$  at  $p \in M$  as a linear map  $df(p): T_p M \rightarrow \mathbb{R}$  by

$$df(p) \cdot v = (f \circ \alpha)'(0),$$

where  $\alpha: (-\epsilon, \epsilon) \rightarrow M$  is a differentiable curve with  $\alpha(0) = p$  and  $\alpha'(0) = v$ .

**Example 1.3.4.** Given a unit vector  $v \in \mathbb{R}^3$ , consider the height function  $f: M \rightarrow \mathbb{R}$  given by  $f(p) = \langle v, p \rangle$ , for every  $p \in M$ . Fix a point  $p \in M$  and let  $w \in T_p M$ . To compute the differential  $df(p) \cdot w$ , choose a differentiable curve  $\alpha: (-\epsilon, \epsilon) \rightarrow M$  with  $\alpha(0) = p$  and  $\alpha'(0) = w$ . Then

$$\begin{aligned} df(p) \cdot w &= (f \circ \alpha)'(0) = \frac{d}{dt} f(\alpha(t))|_{t=0} = \frac{d}{dt} \langle v, \alpha(t) \rangle|_{t=0} \\ &= \langle v, \alpha'(0) \rangle = \langle v, w \rangle. \end{aligned}$$

**Example 1.3.5.** Let  $R_\theta: \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be the rotation of angle  $\theta$  about the axis- $z$ . The rotation  $R_\theta$  is linear, therefore it is differentiable. Moreover,  $R_\theta$  restricted to the unit sphere  $\mathbb{S}^2$  is a differentiable map of  $\mathbb{S}^2$ . Thus, fixed a point  $p \in \mathbb{S}^2$  and given  $v \in T_p \mathbb{S}^2$ , we obtain

$$dR_\theta(p) \cdot v = R_\theta(v),$$

because  $R_\theta$  is linear. Note that  $R_\theta$  leaves the north pole  $N$  fixed, and  $dR_\theta(N)$  is just a rotation of angle  $\theta$  in the plane  $T_N M$ .

## 1.4 Orientable surfaces

Intuitively, orientable surfaces are those for which it is possible to define a clockwise consistently. To illustrate the underlying idea, we consider two familiar surfaces: a cylinder and a Mobius band. We can distinguish between a cylinder and a Mobius band by noticing that every cylinder has an inside and an outside, and we can paint one blue and other yellow, for example. But if we try to paint a Mobius band in two colors, we fail because it has just one side.

Let  $E$  be a finite-dimensional real vector space. We say that two bases  $\mathcal{E}$  and  $\mathcal{F}$  *define the same orientation* in  $E$  if the transition matrix from  $\mathcal{E}$  to  $\mathcal{F}$  has positive determinant. In this case, we write  $\mathcal{E} \equiv \mathcal{F}$ . This property defines an equivalence relation on the set of all bases of  $E$ , and each equivalence class according to this relation is called an *orientation* for  $E$ . Moreover, the relation  $\equiv$  has exactly two equivalence class, that is, the vector space  $E$  admits two orientations.

A vector space  $E$  together with a choice of orientation  $\mathcal{O}$  is called an *oriented vector space*. Once an orientation  $\mathcal{O}$  for  $E$  is fixed, the other one is called the *opposite orientation*, and it will be denoted by  $-\mathcal{O}$ . The bases in the orientation  $\mathcal{O}$  will be called *positive*, while the others will be called *negative*. A linear isomorphism  $T: E \rightarrow F$  between two oriented vector spaces is called *positive* if maps positive bases of  $E$  into positive bases of  $F$ .

**Example 1.4.1.** The Euclidean space  $\mathbb{R}^n$  will be considered oriented requiring that the canonical basis of  $\mathbb{R}^n$  be positive. Therefore, a linear isomorphism  $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$  is positive if and only if  $\det(T) > 0$ .

Now we will extend the notion of orientability to each tangent space of a regular surface  $M$ . We say that two parametrizations  $\varphi: U \rightarrow \varphi(U)$  and  $\psi: V \rightarrow \psi(V)$  of  $M$  are *compatible* if either  $\varphi(U) \cap \psi(V) = \emptyset$  or the change of coordinates

$$\psi^{-1} \circ \varphi: \varphi^{-1}(W) \rightarrow \psi^{-1}(W),$$

has positive jacobian determinant everywhere on  $\varphi^{-1}(W)$ .

**Remark 1.4.2.** If  $\varphi(U) \cap \psi(V) \neq \emptyset$ , the change of coordinates  $\psi^{-1} \circ \varphi$  has jacobian determinant different from zero on  $\varphi^{-1}(\varphi(U) \cap \psi(V))$ . Since determinant is a continuous function, its sign is constant in each connected component of the open set  $\varphi^{-1}(\varphi(U) \cap \psi(V)) \subset \mathbb{R}^2$ .

**Definition 1.4.3.** A regular surface  $M$  is called *orientable* if there exists a cover  $\mathcal{A}$  of  $M$  consisting of coordinate neighborhoods such that any two parameterizations of  $\mathcal{A}$  are compatible.

The choice of such a cover is called an *orientation* of  $M$ , and in this case we say that  $M$  is *oriented*. If it is not possible to make such a choice, the surface  $M$  is called *nonorientable*.

**Example 1.4.4.** The plane  $\mathbb{R}^2$  is an orientable surface, because the identity map is a global compatible parametrization of  $\mathbb{R}^2$ . The orientation given by such parametrization is called the *canonical orientation* of  $\mathbb{R}^2$ .

**Example 1.4.5.** A regular surface which is the graph of a differentiable function is an orientable surface. More generally, all surfaces which can be covered by one coordinate neighborhood are trivially orientable.

**Proposition 1.4.6.** An orientation on a regular surface  $M$  determines an orientation on each tangent plane of  $M$ .

*Proof.* Let  $\mathcal{A}$  be an orientation on  $M$ . Given a point  $p \in M$ , consider a parametrization  $\varphi \in \mathcal{A}$ , with  $p = \varphi(x)$ , and define an orientation  $\mathcal{O}_p$  on  $T_p M$  requiring the basis  $\{d\varphi(x) \cdot e_1, d\varphi(x) \cdot e_2\}$  be positive. If  $\psi$  is another parametrization in the cover  $\mathcal{A}$ , with  $p = \psi(y)$ , we obtain:

$$d\psi(y) = d(\varphi \circ \varphi^{-1} \circ \psi)(y) = d\varphi(x) \circ d(\varphi^{-1} \circ \psi)(y).$$

The isomorphism  $d(\varphi^{-1} \circ \psi)(y)$  preserve orientation, because  $\varphi$  and  $\psi$  are compatible, e  $d\varphi(x)$  preserves orientation by hypothesis. Therefore, the set  $\{d\psi(y) \cdot e_1, d\psi(y) \cdot e_2\}$  is also a positive basis of  $T_p M$ .  $\square$

Before giving a geometric interpretation of the idea of orientability of a regular surface in  $\mathbb{R}^3$ , we need a few definitions.

Given a regular surface  $M$ , the inner product of  $\mathbb{R}^3$  naturally induces an inner product in each tangent plane  $T_p M$  of  $M$ . More precisely, given  $p \in M$  and  $v, w \in T_p M$ , we define  $\langle v, w \rangle$  to be the inner product of  $v$  and  $w$  as vectors of  $\mathbb{R}^3$ .

The inner product defined above allows us to consider the notion of orthogonality. More precisely, we say that a vector  $\eta \in \mathbb{R}^3$  is *orthogonal* to a regular surface  $M$  at a point  $p \in M$  if  $\langle \eta, v \rangle = 0$ , for every  $v \in T_p M$ . Globally, a *normal vector field* to a regular surface  $M$  is a map  $\eta: M \rightarrow \mathbb{R}^3$  such that  $\eta(p)$  is orthogonal to  $T_p M$ , for every  $p \in M$ .

**Theorem 1.4.7.** *A regular surface  $M \subset \mathbb{R}^3$  is orientable if and only if there exists a differentiable unit normal vector field  $N: M \rightarrow \mathbb{R}^3$  on  $M$ .*

*Proof.* Let  $\mathcal{A}$  be an orientation of  $M$ , that is, a cover of  $M$  by compatible coordinate neighborhoods. Fix a point  $p \in M$  with  $p = \varphi(u, v)$ , where  $\varphi: U \rightarrow \varphi(U)$  is a positive parametrization of  $M$ . Now, define a map  $N: \varphi(U) \rightarrow \mathbb{R}^3$  by

$$N(q) = \frac{\varphi_u \times \varphi_v}{\|\varphi_u \times \varphi_v\|}(q), \quad (1.6)$$

for every  $q \in \varphi(U)$ , where  $\times$  denotes the vector product in  $\mathbb{R}^3$ . Thus we obtain a differentiable map, orthogonal to  $M$  in every point of  $\varphi(U)$ . If  $\psi: V \rightarrow \psi(V)$  is another positive parametrization of  $M$  with  $p = \psi(z, w)$ , let us denote by  $h = \varphi^{-1} \circ \psi$  the change of coordinates, with  $(u, v) = h(z, w)$ . Thus  $\psi = \varphi \circ h$  and by setting  $q = (z, w)$ , we obtain

$$\begin{aligned} \psi_z(q) &= \varphi_u(h(q)) \frac{\partial u}{\partial z}(q) + \varphi_v(h(q)) \frac{\partial v}{\partial z}(q) \\ \psi_w(q) &= \varphi_u(h(q)) \frac{\partial u}{\partial w}(q) + \varphi_v(h(q)) \frac{\partial v}{\partial w}(q) \end{aligned}.$$

It follows that

$$\psi_z \times \psi_w = \det(dh(q)) \cdot \varphi_u \times \varphi_v,$$

and from which we conclude that the normals associated to  $\varphi$  and  $\psi$  coincide, since  $\det(dh(q)) > 0$ . Therefore, if  $p$  belongs to the intersection of two coordinate neighborhoods, the corresponding normals coincide, and thus we obtain an unit normal differentiable vector field  $N: M \rightarrow \mathbb{R}^3$ . Conversely, given a point  $p \in M$ , define an orientation in  $T_p M$  as follows: a basis  $\{v_1, v_2\}$  of  $T_p M$  is positive if and only if  $\{v_1, v_2, N(p)\}$  is a positive basis of  $\mathbb{R}^3$ . Given a parametrization  $\varphi: U \rightarrow \varphi(U)$  of  $M$ , with  $p = \varphi(x)$  and  $U$  connected, and changing the sign of  $\varphi$  if necessary, we can assume that the set

$$\{d\varphi(x) \cdot e_1, d\varphi(x) \cdot e_2, N(\varphi(x))\}$$

is a positive basis of  $\mathbb{R}^3$ , for every  $x \in U$ . Thus, for each point  $p \in M$ , we can choose a parametrization  $\varphi: U \rightarrow \varphi(U)$  of  $M$ , with  $p \in \varphi(U)$ , such that  $d\varphi(x): \mathbb{R}^2 \rightarrow T_{\varphi(x)} M$  is a positive linear isomorphism, for every  $x \in U$ . Denote by  $\mathcal{A}$  the cover of  $M$  by such parametrizations. If  $\varphi: U \rightarrow \varphi(U)$  and  $\psi: V \rightarrow \psi(V)$  are two parametrizations in  $\mathcal{A}$ , with  $\varphi(U) \cap \psi(V) \neq \emptyset$ , then  $\psi^{-1} \circ \varphi$  has positive jacobian determinant everywhere, since  $d(\psi^{-1} \circ \varphi)(x)$  is the composite of two positive linear isomorphisms.  $\square$

It follows from the proof of Theorem 1.4.7 that, given a point  $p \in M$  and a parametrization  $\varphi: U \rightarrow \varphi(U)$  of  $M$ , with  $p \in \varphi(U)$ , we can always consider an unit normal differentiable vector field  $N$  in a neighborhood of  $p$  and given by (1.6). Thus, any regular surface  $M$  is always locally orientable.

Let us now look at some examples of orientable surfaces.

**Example 1.4.8.** Let  $\mathcal{P}$  be the plane through the point  $p \in \mathbb{R}^3$  which contains the orthonormal vectors  $w_1, w_2 \in \mathbb{R}^3$ . Thus, a parametrization of  $\mathcal{P}$  is given by

$$\varphi(u, v) = p + uw_1 + vw_2,$$

with  $(u, v) \in \mathbb{R}^2$ . In this case one has  $\varphi_u = w_1$  and  $\varphi_v = w_2$ , therefore  $N = w_1 \times w_2$  is an unit normal vector field along  $\mathcal{P}$ .

**Example 1.4.9.** Consider the right cylinder  $\mathcal{C}$  over the circle  $x^2 + y^2 = 1$ . Then  $\mathcal{C}$  admits a parametrization  $\varphi: U \rightarrow \mathbb{R}^3$  given by

$$\varphi(u, v) = (\cos u \sin u, v),$$

where

$$U = \{(u, v) \in \mathbb{R}^2 : 0 < u < 2\pi \text{ and } v \in \mathbb{R}\}.$$

In this case, we obtain

$$\varphi_u = (-\sin u, \cos u, 0) \quad \text{and} \quad \varphi_v = (0, 0, 1),$$

which implies that

$$N = \varphi_u \times \varphi_v = (\cos u, \sin u, 0)$$

is an unit normal vector field to the cylinder  $\mathcal{C}$ .

**Example 1.4.10.** The tangent plane to the unit sphere  $\mathbb{S}^2 \subset \mathbb{R}^3$  at a point  $p \in \mathbb{S}^2$  is given by

$$T_p \mathbb{S}^2 = p^\perp,$$

where  $p^\perp = \{v \in \mathbb{R}^3 : \langle v, p \rangle = 0\}$ . In fact, fix a vector  $v \in T_p \mathbb{S}^2$  with  $v = \alpha'(0)$ , where  $\alpha: (-\epsilon, \epsilon) \rightarrow \mathbb{S}^2$  is a differentiable curve, with  $\alpha(0) = p$ . Since  $\alpha(t) \in \mathbb{S}^2$  for every  $t \in (-\epsilon, \epsilon)$ , one has  $\|\alpha(t)\| = 1$ , for every  $t \in (-\epsilon, \epsilon)$ . This implies that

$$2\langle \alpha'(t), \alpha(t) \rangle = 0,$$

for every  $t \in (-\epsilon, \epsilon)$ . For  $t = 0$ , we obtain  $\langle v, p \rangle = 0$ . This shows that  $T_p \mathbb{S}^2 \subset p^\perp$  and hence the equality by dimensional reasons. Therefore, the position vector field  $N: \mathbb{S}^2 \rightarrow \mathbb{R}^3$ ,  $N(p) = p$ , is an unit normal vector field to the sphere, and this shows that  $\mathbb{S}^2$  is orientable.

**Example 1.4.11.** Let  $M$  be a regular surface given as the inverse image under a differentiable function  $f: \mathbb{R}^3 \rightarrow \mathbb{R}$  of a regular value  $c \in \mathbb{R}$ , that is,  $M = f^{-1}(c)$ . Fix a point  $p \in M$ , with  $p = (x_0, y_0, z_0)$ , and consider a differentiable parametrized curve  $\alpha: (-\epsilon, \epsilon) \rightarrow M$ , with  $\alpha(0) = p$  and

$$\alpha(t) = (x(t), y(t), z(t)).$$

Since  $\alpha(t) \in M$  for every  $t \in (-\epsilon, \epsilon)$ , we obtain

$$f(\alpha(t)) = c, \tag{1.7}$$

for every  $t \in (-\epsilon, \epsilon)$ . By differentiating both sides of (1.7) with respect to  $t$ , we see that at  $t = 0$

$$\begin{aligned} 0 &= \frac{d}{dt} f(\alpha(t))(0) = \frac{\partial f}{\partial x}(p) \cdot \frac{dx}{dt}(0) + \frac{\partial f}{\partial y}(p) \cdot \frac{dy}{dt}(0) + \frac{\partial f}{\partial z}(p) \cdot \frac{dz}{dt}(0) \\ &= \left\langle \left( \frac{\partial f}{\partial x}(p), \frac{\partial f}{\partial y}(p), \frac{\partial f}{\partial z}(p) \right), \left( \frac{dx}{dt}(0), \frac{dy}{dt}(0), \frac{dz}{dt}(0) \right) \right\rangle \\ &= \langle \text{grad} f(p), \alpha'(0) \rangle = \langle \text{grad} f(p), v \rangle. \end{aligned}$$

This shows that the gradient vector of  $f$  at  $p \in M$  is orthogonal to  $T_p M$ . Therefore, the map

$$N(p) = \frac{\text{grad} f}{\|\text{grad} f\|}(p)$$

is a unit normal vector field to  $M$ , and thus  $M$  is orientable.

## Chapter 2

# The local theory of surfaces

### 2.1 Derivatives of vector fields in $\mathbb{R}^3$

A *vector field* defined in an open set  $U \subset \mathbb{R}^3$  is simply a differentiable map  $X: U \rightarrow \mathbb{R}^3$  that maps each point  $p \in U$  in a vector  $X(p) \in \mathbb{R}^3$ . The set of all vector fields defined in the open  $U \subset \mathbb{R}^3$  will be denoted by  $\mathfrak{X}(U)$ . Under the natural operations

$$\begin{aligned}(X + Y)(p) &= X(p) + Y(p) \\ (c \cdot X)(p) &= c \cdot X(p)\end{aligned},$$

this set  $\mathfrak{X}(U)$  is a real vector space. The next question we want to address is to define a notion of derivative of vector fields along a vector, and more generally, along another vector field.

Let  $Y \in \mathfrak{X}(U)$  be a vector field and fix a point  $p \in U$ . The *directional derivative* of  $Y$  at  $p$ , along a vector  $v \in \mathbb{R}^3$ , is defined by

$$\tilde{\nabla}_v Y = dY(p) \cdot v. \quad (2.1)$$

Note that if  $\alpha: (-\epsilon, \epsilon) \rightarrow U$  is a differentiable curve with  $\alpha(0) = p$  and  $\alpha'(0) = v$ , then

$$\begin{aligned}\lim_{t \rightarrow 0} \frac{Y(\alpha(t)) - Y(\alpha(0))}{t} &= \frac{d}{dt} Y(\alpha(t))|_{t=0} \\ &= dY(p) \cdot v \\ &= \tilde{\nabla}_v Y.\end{aligned}$$

This implies that the vector  $\tilde{\nabla}_v Y$  depends only on the values of  $Y$  along a differentiable curve through  $p$  with velocity  $v$ . We can extend the above

definition. More precisely, given two vector fields  $X, Y \in \mathfrak{X}(U)$ , we will define another vector field  $\tilde{\nabla}_X Y$  by

$$\left(\tilde{\nabla}_X Y\right)(p) = \tilde{\nabla}_{X(p)} Y. \quad (2.2)$$

In the same way as in (2.1), the left-hand side of (2.2) depends only on the values of  $Y$  along a differentiable curve through  $p$  with velocity  $X(p)$ .

To understand more fully what the derivative in (2.1) means, it is useful to bear in mind some terminology. Fix a point  $p \in U$ . Given a vector  $v \in \mathbb{R}^3$ , we can define a natural function  $v: C^\infty(U) \rightarrow \mathbb{R}$  by

$$v(f) = \frac{d}{dt} f(\alpha(t))|_{t=0} = \frac{\partial f}{\partial v}(p) = \sum_{i=1}^3 \frac{\partial f}{\partial x_i}(p) \cdot v_i, \quad (2.3)$$

for every  $f \in C^\infty(U)$ , where  $\alpha: (-\epsilon, \epsilon) \rightarrow U$  is a differentiable curve with  $\alpha(0) = p$  and  $\alpha'(0) = v$ , and  $v = (v_1, v_2, v_3)$ . The function given in (2.3) is called a *derivation* at  $p$  along the vector  $v$ . On the other hand, given a vector field  $X \in \mathfrak{X}(U)$  and a function  $f \in C^\infty(U)$ , we can define another function  $X(f): U \rightarrow \mathbb{R}$  by

$$X(f)(p) = X(p)(f) = df(p) \cdot X(p),$$

where  $X(p)(f)$  is the derivation at  $p$  along the vector  $X(p)$  given in (2.3).

Therefore, motivated by these considerations, fix a point  $p \in U$  and given a vector field  $Y \in \mathfrak{X}(U)$ , with  $Y = (Y_1, Y_2, Y_3)$ , we can write

$$\begin{aligned} \tilde{\nabla}_v Y &= \frac{d}{dt} Y(\alpha(t))|_{t=0} = \frac{d}{dt} (Y_1(\alpha(t)), Y_2(\alpha(t)), Y_3(\alpha(t)))|_{t=0} \\ &= (v(Y_1), v(Y_2), v(Y_3)), \end{aligned} \quad (2.4)$$

where  $\alpha: (-\epsilon, \epsilon) \rightarrow U$  is a differentiable curve with  $\alpha(0) = p$  and  $\alpha'(0) = v$ . In other words, the directional derivative of a vector field along a vector is just a derivation of its coordinates functions related to that vector.

In the sequel, we will define a map

$$\tilde{\nabla}: \mathfrak{X}(U) \times \mathfrak{X}(U) \rightarrow \mathfrak{X}(U) \quad (2.5)$$

that assigns to each pair  $X, Y \in \mathfrak{X}(U)$  the vector field  $\tilde{\nabla}_X Y \in \mathfrak{X}(U)$  given in (2.2). The map  $\tilde{\nabla}$  defined in (2.5) is called the *derivative* of the vector fields in the open set  $U \subset \mathbb{R}^3$ .

**Proposition 2.1.1.** The map  $\tilde{\nabla}$  satisfies the following basic properties:

- (a)  $\tilde{\nabla}_{fX}Y = f\tilde{\nabla}_XY,$
- (b)  $\tilde{\nabla}_XfY = X(f)Y + f\tilde{\nabla}_XY,$

for every  $X, Y \in \mathfrak{X}(U)$  and  $f \in C^\infty(U)$ .

*Proof.* Fix a point  $p \in U$ . By writing  $Y = (Y_1, Y_2, Y_3)$ , we obtain

$$\begin{aligned} \left(\tilde{\nabla}_{fX}Y\right)(p) &= (f(p)X_pY_1, f(p)X_pY_2, f(p)X_pY_3) \\ &= f(p)(X_pY_1, X_pY_2, X_pY_3) \\ &= f(p)\left(\tilde{\nabla}_XY\right)(p) \end{aligned}$$

and

$$\begin{aligned} \left(\tilde{\nabla}_XfY\right)(p) &= (X_p(fY_1), X_p(fY_2), X_p(fY_3)) \\ &= (X(f)Y_1 + fX(Y_1), \dots, X(f)Y_3 + fX(Y_3))(p) \\ &= X_p(f)(Y_1, Y_2, Y_3) + f(p)\tilde{\nabla}_{X_p}Y \\ &= \left(X(f)Y + f\tilde{\nabla}_XY\right)(p), \end{aligned}$$

and this concludes the proof.  $\square$

**Proposition 2.1.2.** Given  $X, Y \in \mathfrak{X}(U)$ , there exists a unique vector field  $Z \in \mathfrak{X}(U)$  such that

$$Z(f) = (XY - YX)(f), \quad (2.6)$$

for every  $f \in C^\infty(U)$ .

*Proof.* Writing  $X = (X_1, X_2, X_3)$  and  $Y = (Y_1, Y_2, Y_3)$ , we obtain

$$XY(f) = X\left(\sum_{j=1}^3 Y_j \frac{\partial f}{\partial x_j}\right) = \sum_{i,j=1}^3 \left(X_i \frac{\partial Y_j}{\partial x_i} \frac{\partial f}{\partial x_j} + X_i Y_j \frac{\partial^2 f}{\partial x_i \partial x_j}\right)$$

and

$$YX(f) = Y\left(\sum_{i=1}^3 X_i \frac{\partial f}{\partial x_i}\right) = \sum_{i,j=1}^3 \left(Y_j \frac{\partial X_i}{\partial x_j} \frac{\partial f}{\partial x_i} + X_i Y_j \frac{\partial^2 f}{\partial x_j \partial x_i}\right).$$

Thus,

$$(XY - YX)(f) = \sum_{i,j=1}^3 \left( X_i \frac{\partial Y_j}{\partial x_i} - Y_i \frac{\partial X_j}{\partial x_i} \right) \frac{\partial f}{\partial x_j}. \quad (2.7)$$

Therefore, by defining the vector field  $Z = (Z_1, Z_2, Z_3)$  by

$$Z_j = \sum_{i=1}^3 \left( X_i \frac{\partial Y_j}{\partial x_i} - Y_i \frac{\partial X_j}{\partial x_i} \right),$$

it follows that  $Z$  is the unique vector field in  $U$  satisfying (2.6).  $\square$

The vector field  $Z$ , given by Proposition 2.1.2, is called the *Lie bracket* of  $X$  and  $Y$ , and it will be denoted by  $[X, Y]$ .

**Proposition 2.1.3.** The map  $\tilde{\nabla}$  satisfies the following properties:

- (a)  $X\langle Y, Z \rangle = \langle \tilde{\nabla}_X Y, Z \rangle + \langle Y, \tilde{\nabla}_X Z \rangle$ ,
- (b)  $\tilde{\nabla}_X Y - \tilde{\nabla}_Y X = [X, Y]$ ,

for every  $X, Y, Z \in \mathfrak{X}(U)$ . Conversely, if

$$\overline{\nabla} : \mathfrak{X}(U) \times \mathfrak{X}(U) \rightarrow \mathfrak{X}(U)$$

is a map that satisfies the properties (a) and (b) above, then  $\overline{\nabla} = \tilde{\nabla}$ .

*Proof.* Part (a) follows from

$$\begin{aligned} X\langle Y, Z \rangle(p) &= X(p) \left( \sum_{i=1}^3 Y_i Z_i \right) \\ &= \sum_{i=1}^3 (X(p)(Y_i) Z_i + Y_i X(p)(Z_i)) \\ &= \langle \tilde{\nabla}_{X(p)} Y, Z(p) \rangle + \langle Y(p), \tilde{\nabla}_{X(p)} Z \rangle, \end{aligned}$$

for every  $p \in U$ . For part (b), writing  $\tilde{\nabla}_X Y - \tilde{\nabla}_Y X = (W_1, W_2, W_3)$  and using (2.4), we obtain

$$W_i = X(p)(Y_i) - Y(p)(X_i) = \sum_{j=1}^3 \left( X_j \frac{\partial Y_i}{\partial x_j}(p) - Y_j \frac{\partial X_i}{\partial x_j}(p) \right),$$

and the formula (b) follows from (2.7). Conversely, suppose that a map  $\bar{\nabla} : \mathfrak{X}(U) \times \mathfrak{X}(U) \rightarrow \mathfrak{X}(U)$  satisfies the properties (a) and (b). By part (a), one has

$$X\langle Y, Z \rangle = \langle \bar{\nabla}_X Y, Z \rangle + \langle Y, \bar{\nabla}_X Z \rangle \quad (2.8)$$

$$Y\langle X, Z \rangle = \langle \bar{\nabla}_Y X, Z \rangle + \langle X, \bar{\nabla}_Y Z \rangle \quad (2.9)$$

$$Z\langle X, Y \rangle = \langle \bar{\nabla}_Z X, Y \rangle + \langle X, \bar{\nabla}_Z Y \rangle, \quad (2.10)$$

for every  $X, Y, Z \in \mathfrak{X}(U)$ . Adding (2.8) and (2.9), subtracting (2.10) and using part (b), we obtain

$$\begin{aligned} X\langle Y, Z \rangle + Y\langle X, Z \rangle - Z\langle X, Y \rangle &= \langle [X, Z], Y \rangle + \langle [Y, Z], X \rangle \\ &\quad + \langle [X, Y], Z \rangle + 2\langle \bar{\nabla}_Y X, Z \rangle \end{aligned}$$

that is,

$$\begin{aligned} 2\langle \bar{\nabla}_Y X, Z \rangle &= X\langle Y, Z \rangle + Y\langle X, Z \rangle - Z\langle X, Y \rangle - \langle [X, Z], Y \rangle \\ &\quad - \langle [Y, Z], X \rangle - \langle [X, Y], Z \rangle \end{aligned} \quad (2.11)$$

The equation (2.11) shows that  $\bar{\nabla}$  is uniquely determined by the inner product  $\langle \cdot, \cdot \rangle$ , and therefore

$$\langle \bar{\nabla}_Y X, Z \rangle = \langle \tilde{\nabla}_Y X, Z \rangle,$$

for every  $X, Y, Z \in \mathfrak{X}(U)$ , and this shows that  $\bar{\nabla} = \tilde{\nabla}$ .  $\square$

**Proposition 2.1.4.** For any vector fields  $X, Y, Z \in \mathfrak{X}(U)$ , the following holds:

$$\tilde{\nabla}_X \tilde{\nabla}_Y Z - \tilde{\nabla}_Y \tilde{\nabla}_X Z - \tilde{\nabla}_{[X, Y]} Z = 0. \quad (2.12)$$

*Proof.* By definition, we have

$$\tilde{\nabla}_X \tilde{\nabla}_Y Z = \tilde{\nabla}_X (Y Z_1, Y Z_2, Y Z_3) = (XY Z_1, XY Z_2, XY Z_3)$$

and

$$\tilde{\nabla}_Y \tilde{\nabla}_X Z = \tilde{\nabla}_Y (X Z_1, X Z_2, X Z_3) = (YX Z_1, YX Z_2, YX Z_3).$$

This implies that

$$\begin{aligned} \tilde{\nabla}_X \tilde{\nabla}_Y Z - \tilde{\nabla}_Y \tilde{\nabla}_X Z &= (XY Z_1 - YX Z_1, XY Z_2 - YX Z_2, XY Z_3 - YX Z_3) \\ &= ([X, Y] Z_1, [X, Y] Z_2, [X, Y] Z_3) \\ &= \tilde{\nabla}_{[X, Y]} Z, \end{aligned}$$

and this proves equation (2.12).  $\square$

## 2.2 Derivatives of vector fields on surfaces

A *vector field along* a regular surface  $M$  in  $\mathbb{R}^3$  is just a differentiable map  $X: M \rightarrow \mathbb{R}^3$  that maps each point  $p \in M$  in a vector  $X(p) \in \mathbb{R}^3$ . We say that  $X$  is *tangent* to  $M$  if  $X(p) \in T_p M$  for every  $p \in M$ . We also say that  $X$  is *normal* to  $M$  if  $X(p) \perp T_p M$  for every  $p \in M$ . We denote by  $\mathfrak{X}(M)$  the set of all tangent vector fields to  $M$ . In a similar way to the case of vector fields in  $\mathbb{R}^3$ , the set  $\mathfrak{X}(M)$  becomes a real vector space endowed with the natural operations.

Given two vector fields  $X, Y \in \mathfrak{X}(M)$ , we define a map  $[X, Y]: M \rightarrow \mathbb{R}^3$  by

$$[X, Y](p) = [\tilde{X}, \tilde{Y}](p) \quad (2.13)$$

for every  $p \in M$ , where  $\tilde{X}, \tilde{Y}$  are differentiable extensions of  $X$  and  $Y$ , respectively, to an open set  $U \subset \mathbb{R}^3$ , with  $p \in U$ . We will see that (2.13) is well defined and  $[X, Y] \in \mathfrak{X}(M)$  for any choice of vector fields  $X, Y \in \mathfrak{X}(M)$ .

Given a parametrization  $\varphi: U \rightarrow \varphi(U)$  of  $M$ , we can write on the coordinate neighborhood  $\varphi(U)$

$$X = \sum_{i=1}^2 a_i \frac{\partial}{\partial x_i} \quad \text{and} \quad Y = \sum_{j=1}^2 b_j \frac{\partial}{\partial x_j},$$

where  $\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}$  are the coordinate vector fields associated to  $\varphi$ .

**Proposition 2.2.1.** On the coordinate neighborhood  $\varphi(U)$ , we have

$$[X, Y] = \sum_{i,j=1}^2 \left( a_i \frac{\partial b_j}{\partial x_i} - b_i \frac{\partial a_j}{\partial x_i} \right) \frac{\partial}{\partial x_j}. \quad (2.14)$$

*Proof.* Consider the map  $\phi: U \times (-\epsilon, \epsilon) \rightarrow \mathbb{R}^3$  given by

$$\phi(x_1, x_2, x_3) = \varphi(x_1, x_2) + x_3 N(\varphi(x_1, x_2)),$$

where  $N$  is a unit normal vector field to  $M$  along  $\varphi(U)$ . Note that  $\varphi$  is differentiable and

$$\begin{aligned} \frac{\partial \phi}{\partial x_1} &= \frac{\partial \varphi}{\partial x_1} + x_3 \frac{\partial N}{\partial x_1}, \\ \frac{\partial \phi}{\partial x_2} &= \frac{\partial \varphi}{\partial x_2} + x_3 \frac{\partial N}{\partial x_2}, \\ \frac{\partial \phi}{\partial x_3} &= N. \end{aligned}$$

Since  $\frac{\partial \varphi}{\partial x_1}, \frac{\partial \varphi}{\partial x_2}, N$  are linearly independent, the same is true for  $\frac{\partial \phi}{\partial x_1}, \frac{\partial \phi}{\partial x_2}, \frac{\partial \phi}{\partial x_3}$ , for sufficiently small  $\epsilon > 0$ . Restricting the range  $U \times (-\epsilon, \epsilon)$  if necessary, it follows from the inverse function theorem that  $\phi$  is a diffeomorphism between  $U \times (-\epsilon, \epsilon)$  and  $\phi(U \times (-\epsilon, \epsilon))$ . Thus, if

$$\tilde{X} = \sum_{i=1}^3 a_i \frac{\partial}{\partial x_i} \quad \text{and} \quad \tilde{Y} = \sum_{j=1}^3 b_j \frac{\partial}{\partial x_j}$$

are differentiable extensions of  $X|_{\varphi(U)}$  and  $Y|_{\varphi(U)}$ , respectively, it follows from (2.7) that

$$[\tilde{X}, \tilde{Y}] = \sum_{i,j=1}^3 \left( a_i \frac{\partial b_j}{\partial x_i} - b_i \frac{\partial a_j}{\partial x_i} \right) \frac{\partial}{\partial x_j}.$$

Therefore,

$$[\tilde{X}, \tilde{Y}]|_{\varphi(U)} = \sum_{i,j=1}^3 \left( a_i \frac{\partial b_j}{\partial x_i} - b_i \frac{\partial a_j}{\partial x_i} \right) \frac{\partial}{\partial x_j},$$

and this concludes the proof.  $\square$

Similarly as vector fields in  $\mathbb{R}^3$ , the vector field  $[X, Y] \in \mathfrak{X}(M)$  will be called the *Lie bracket* of  $X$  and  $Y$ . It follows directly from (2.14) the following

**Corollary 2.2.2.** If  $\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}$  are the coordinate vector fields associated to a parametrization  $\varphi$  of  $M$ , then

$$\left[ \frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2} \right] = 0.$$

**Example 2.2.3.** In  $\mathbb{R}^2$ , with canonical coordinates  $(x, y)$ , consider the vector fields  $X = y \frac{\partial}{\partial y}$  and  $Y = x \frac{\partial}{\partial y}$ . Given a function  $f \in C^\infty(\mathbb{R}^2)$ , we have

$$\begin{aligned} [X, Y](f) &= \left[ y \frac{\partial}{\partial y}, x \frac{\partial}{\partial y} \right] (f) \\ &= yx \frac{\partial^2 f}{\partial y^2} - x \frac{\partial f}{\partial y} - xy \frac{\partial^2 f}{\partial y^2} \\ &= -x \frac{\partial}{\partial y} (f) = -Y(f). \end{aligned}$$

Therefore, in this case one has  $[X, Y] = -Y$ .

Let  $f: M \rightarrow N$  be a diffeomorphism between two regular surfaces  $M$  and  $N$ . Given a vector field  $X \in \mathfrak{X}(M)$ , denote by  $f_*X$  the vector field in  $N$  given by

$$(f_*X)(f(p)) = df(p) \cdot X(p),$$

for every  $p \in M$ .

**Corollary 2.2.4.** If  $f: M \rightarrow N$  is a diffeomorphism, then

$$[f_*X, f_*Y] = f_*[X, Y], \quad (2.15)$$

for all  $X, Y \in \mathfrak{X}(M)$ .

*Proof.* Given a parametrization  $\varphi: U \rightarrow \varphi(U)$  of  $M$ , we can write  $X$  and  $Y$  along  $\varphi(U)$  as

$$X = \sum_{i=1}^2 a_i \frac{\partial}{\partial x_i} \quad \text{and} \quad Y = \sum_{j=1}^2 b_j \frac{\partial}{\partial x_j}.$$

Let  $\psi: U \rightarrow \psi(V)$  be the map given  $\psi = f \circ \varphi$ , where  $V = f(\varphi(U))$ . Since  $\varphi$  is a parametrization of  $M$  and  $f$  is a diffeomorphism, it follows that  $\psi$  is a parametrization of  $N$ , and the coordinate vector fields  $\frac{\partial}{\partial y_1}, \frac{\partial}{\partial y_2}$  associated to  $\psi$  are given by

$$\frac{\partial}{\partial y_i} = \psi_* \frac{\partial}{\partial x_i}, \quad 1 \leq i \leq 2.$$

Thus, along  $\psi(V)$  we have

$$\psi_*X = \sum_{i=1}^2 \tilde{a}_i \frac{\partial}{\partial y_i} \quad \text{and} \quad \psi_*Y = \sum_{j=1}^2 \tilde{b}_j \frac{\partial}{\partial y_j},$$

where  $a_i \circ \psi = \tilde{a}_i$  and  $b_j \circ \psi = \tilde{b}_j$ , and equation (2.15) follows from (2.14).  $\square$

Given a tangent vector field  $X \in \mathfrak{X}(M)$  and a vector field  $Y$  along  $M$ , define

$$\left( \tilde{\nabla}_X Y \right) (p) = \left( \tilde{\nabla}_{\tilde{X}} \tilde{Y} \right) (p), \quad (2.16)$$

for every  $p \in M$ , where  $\tilde{X}, \tilde{Y}$  are differentiable extensions of  $X$  and  $Y$ , respectively. Since the right hand side of (2.16) does not depend on the choice of the extensions  $\tilde{X}$  and  $\tilde{Y}$ , the vector field  $\tilde{\nabla}_X Y$  is well defined. Given  $X, Y \in \mathfrak{X}(M)$ , we define a vector field  $\nabla_X Y$  by

$$(\nabla_X Y)(p) = \left( \tilde{\nabla}_X Y \right)^T (p), \quad (2.17)$$

where the right hand side of (2.17) denotes the tangent component of  $\tilde{\nabla}_X Y$  along the surface  $M$ .

**Proposition 2.2.5.** The map  $\nabla: \mathfrak{X}(M) \times \mathfrak{X}(M) \rightarrow \mathfrak{X}(M)$  given in (2.17) satisfies the following properties:

- (a)  $\nabla_{fX}Y = f\nabla_XY$ ,
- (b)  $\nabla_XfY = X(f)Y + f\nabla_XY$ ,
- (c)  $X\langle Y, Z \rangle = \langle \nabla_XY, Z \rangle + \langle Y, \nabla_XZ \rangle$ ,
- (d)  $\nabla_XY - \nabla_YZ = [X, Y]$ ,

for all  $X, Y, Z \in \mathfrak{X}(M)$  and every  $f \in C^\infty(N)$ . Conversely, if a map  $\overline{\nabla}: \mathfrak{X}(M) \times \mathfrak{X}(M) \rightarrow \mathfrak{X}(M)$  satisfies properties (a)–(d) above, then  $\overline{\nabla} = \nabla$ .

*Proof.* Parts (a) and (b) follow by taking the tangent component of formulas (a) and (b), respectively, of Proposition 2.1.1. Analogously, part (c) follows by taking the tangent component of formula (c) of Proposition 2.1.3. Part (d) follows directly of formula (d) of Proposition 2.1.3. The converse is entirely analogous to Proposition 2.1.3.  $\square$

**Corollary 2.2.6.** Let  $f: M \rightarrow N$  be a local isometry between two regular surfaces  $M$  and  $N$ . Then

$$f_*(\nabla_XY) = \nabla_{f_*X}f_*Y,$$

for every  $X, Y \in \mathfrak{X}(M)$ .

*Proof.* Define a map  $\overline{\nabla}: \mathfrak{X}(M) \times \mathfrak{X}(M) \rightarrow \mathfrak{X}(M)$  by

$$\overline{\nabla}_XY = f_*^{-1}(\nabla_{f_*X}f_*Y).$$

A straightforward computation shows that  $\overline{\nabla}$  satisfies properties (a)–(d) of Proposition 2.2.5. For example, property (d) follows from

$$\begin{aligned} \overline{\nabla}_XY - \overline{\nabla}_YX &= f_*^{-1}(\nabla_{f_*X}f_*Y - \nabla_{f_*Y}f_*X) \\ &= f_*^{-1}([f_*X, f_*Y]) \\ &= f_*^{-1}(f_*[X, Y]) \\ &= [X, Y], \end{aligned}$$

using Corollary 2.2.4. Therefore, by the uniqueness of derivative it follows that  $\overline{\nabla}$  coincides with the derivative  $\nabla$  of  $M$ .  $\square$

## 2.3 The Gauss map of a regular surface

Let  $M$  be an orientable regular surface. This means, according to Theorem 1.4.7, that there exists a differentiable unit vector field  $N: M \rightarrow \mathbb{R}^3$ , which is normal along  $M$  and called an *orientation* for  $M$ . Fixed an orientation  $N$  of  $M$ , note that the map  $N: M \rightarrow \mathbb{R}^3$  maps each point  $p \in M$  into a vector  $N(p)$  in the unit sphere  $\mathbb{S}^2 \subset \mathbb{R}^3$ . The map

$$N: M \rightarrow \mathbb{S}^2,$$

thus defined, is called the *Gauss map* of the surface  $M$ .

The Gauss map is differentiable and its differential  $dN(p)$  at a point  $p \in M$  is a linear map from  $T_p M$  to  $T_{N(p)} \mathbb{S}^2$ . Since the planes  $T_p M$  and  $T_{N(p)} \mathbb{S}^2$  are parallel, because they have the same normal vector  $N(p)$ , we can identify them and, thus,  $dN(p)$  is a linear operator on  $T_p M$ .

From a variational point of view, the differential  $dN(p)$  measures the rate of change of the normal vector  $N$  in relation to the tangent plane  $T_p M$  in a neighborhood of  $p$ . In fact, given a vector  $v \in T_p M$ , let  $\lambda: (-\epsilon, \epsilon) \rightarrow M$  be a differentiable curve, with  $\lambda(0) = p$  and  $\lambda'(0) = v$ , and consider the restriction  $N(t) = N(\lambda(t))$ , which is a differentiable curve in the sphere  $\mathbb{S}^2$ . Since  $\|N\| = 1$ , the vector  $N'(0) = dN(p) \cdot v$  is tangent to  $M$  at  $p$ , and it measures the rate of change of  $N$ , restricted to the curve  $\lambda$ , at  $t = 0$ .

We will make some considerations in order to obtain a geometric interpretation for the differential of the Gauss map. Given two vector fields  $X, Y \in \mathfrak{X}(M)$ , define

$$\alpha(X, Y) = \left( \tilde{\nabla}_X Y \right)^\perp, \quad (2.18)$$

where the right-hand side of (2.18) denotes the normal component of  $\tilde{\nabla}_X Y$  along  $M$ . Taking normal components in the equation

$$\tilde{\nabla}_X Y - \tilde{\nabla}_Y X = [X, Y],$$

and using the fact that  $[X, Y] \in \mathfrak{X}(M)$ , we obtain

$$\alpha(X, Y) = \alpha(Y, X),$$

for every  $X, Y \in \mathfrak{X}(M)$ . Since  $\left( \tilde{\nabla}_X Y \right)^\perp(p)$  depends only of  $X(p)$ , it follows from the symmetry of  $\alpha$  that  $\alpha(X, Y)(p)$  depends only of  $X(p)$  e  $Y(p)$ . Thus, for each point  $p \in M$ , we obtain a symmetric bilinear map

$$\alpha_p: T_p M \times T_p M \rightarrow T_p M^\perp$$

given by  $\alpha_p(X_p, Y_p) = \alpha(X, Y)(p)$ . The bilinear map  $\alpha_p$  is usually called the *second fundamental form* of  $M$  at  $p$ . Such map defines a unique linear operator  $A: T_p M \rightarrow T_p M$  by the relation

$$\langle AX, Y \rangle = \langle \alpha_p(X, Y), N(p) \rangle,$$

for every  $X, Y \in T_p M$ . The symmetry of  $\alpha$  implies that  $A$  is a self-adjoint linear operator, and it will be called the *shape operator* of  $M$  at  $p$ .

**Proposition 2.3.1.** Fixed a point  $p \in M$ , we have

$$AX = -\tilde{\nabla}_X N,$$

for every  $X \in T_p M$ .

*Proof.* Given  $X, Y \in T_p M$ , one has

$$\begin{aligned} \langle AX, Y \rangle &= \langle \alpha_p(X, Y), N(p) \rangle \\ &= \left\langle \tilde{\nabla}_X Y, N(p) \right\rangle \\ &= -\left\langle \tilde{\nabla}_X N, Y \right\rangle. \end{aligned}$$

Since  $Y \in T_p M$  is arbitrary, this completes the proof.  $\square$

From Proposition 2.3.1, we identify the shape operator  $A: T_p M \rightarrow T_p M$  with the differential  $dN(p)$  of the Gauss map at the point  $p \in M$ . Since  $A$  is self-adjoint, it follows from the spectral theorem that there exists an orthonormal basis  $\{e_1, e_2\}$  of  $T_p M$  such that

$$Ae_i = k_i e_i, \quad 1 \leq i \leq 2. \quad (2.19)$$

Moreover, the real numbers  $k_1$  and  $k_2$  ( $k_1 \geq k_2$ ) are the maximum and minimum, respectively, of the quadratic form  $\Pi_p$  defined in  $T_p M$  by

$$\Pi_p(v) = -\langle Av, v \rangle, \quad (2.20)$$

restricted to the unit circle of  $T_p M$ .

**Example 2.3.2.** Consider a plane  $\mathcal{P}$  in  $\mathbb{R}^3$  given by  $ax + by + cz + d = 0$ . An unit vector field  $N$  normal to  $\mathcal{P}$  is given by

$$N = \frac{(a, b, c)}{\sqrt{a^2 + b^2 + c^2}},$$

which is constant. This implies that the shape operator  $A = -dN(p)$  vanishes identically at any point  $p$ .

**Example 2.3.3.** In the unit sphere  $\mathbb{S}^2$ , we fix an orientation by choosing  $N(p) = -p$  as a normal vector field. Thus, the shape operator is

$$Av = -dN(p) \cdot v = v,$$

for every  $v \in T_p\mathbb{S}^2$ .

**Example 2.3.4.** In the cylinder

$$\mathcal{C} = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 = 1\},$$

we fix an orientation by choosing  $N = (-x, -y, 0)$  as a normal vector field. Thus, given  $p \in \mathcal{C}$  and  $v \in T_p\mathcal{C}$ , with  $v = (v_1, v_2, v_3)$ , one has

$$Av = -dN(p) \cdot v = (v_1, v_2, 0).$$

In particular, if  $v \in T_p\mathcal{C}$  is parallel to the axis- $z$ , then  $Av = 0 \cdot v$ ; if  $w \in T_p\mathcal{C}$  is parallel to the plane- $xy$ , then  $Aw = 1 \cdot w$ . It follows that the vectors  $v$  and  $w$  are eigenvectors of  $A$  with eigenvalues 0 and 1, respectively.

**Example 2.3.5.** A parametrization for the hyperbolic paraboloid

$$\mathcal{H} = \{(x, y, z) \in \mathbb{R}^3 : z = y^2 - x^2\}$$

is given by

$$\varphi(x, y) = (x, y, y^2 - x^2).$$

Thus, the coordinate vector fields are

$$\frac{\partial \varphi}{\partial x} = (1, 0, -2x) \quad \text{and} \quad \frac{\partial \varphi}{\partial y} = (0, 1, 2y),$$

and an unit vector field  $N$  normal to  $\mathcal{H}$  is

$$N(x, y) = \frac{1}{\sqrt{x^2 + y^2 + 1/4}}(x, -y, 1/2). \quad (2.21)$$

At the origem  $p = (0, 0, 0)$ , the tangent plane  $T_p\mathcal{H}$  coincides with the plane- $xy$ , since  $\frac{\partial \varphi}{\partial x}, \frac{\partial \varphi}{\partial y}$  agree with the unit canonical vectors fields  $e_1$  and  $e_2$ , respectively. Given  $v \in T_p\mathcal{H}$ , let  $\lambda: (-\epsilon, \epsilon) \rightarrow \mathcal{H}$  be a differentiable curve with  $\lambda(0) = p$  and  $\lambda'(0) = v$ . If  $\lambda(t) = (x(t), y(t), z(t))$ , then

$$\lambda'(0) = (x'(0), y'(0), 0).$$

If  $N(t) = N(\lambda(t))$ , it follows from (2.21) that

$$N'(0) = 2(x'(0), -y'(0), 0).$$

Thus, at the origem  $p = (0, 0, 0)$ , the shape operator  $A$  is given by

$$Av = 2(-v_1, v_2, 0),$$

where  $v = (v_1, v_2, v_3)$ . This implies that the canonical vectors  $e_1 = (1, 0, 0)$  and  $e_2 = (0, 1, 0)$  are eigenvectors of  $A$ , with eigenvalues  $-2$  and  $2$ , respectively.

## 2.4 Euler's principal curvatures

Let  $M$  be a regular surface in  $\mathbb{R}^3$  and fix a point  $p \in M$ . Given a regular curve  $\gamma: (-\epsilon, \epsilon) \rightarrow M$ , with  $\gamma(0) = p$ , denote by  $\cos \theta = \langle n(0), N(p) \rangle$  the angle between the normal vector  $n$  to  $\gamma$  and the normal vector  $N$  to  $M$  at  $p$ . The number

$$k_n = k \cos \theta$$

is called the *normal curvature* of  $\gamma$  at  $p \in M$ , where  $k$  is the curvature of  $\gamma$  at  $p$ . Note that  $k_n$  is the length of the projection of the vector  $kn$  over the normal  $N$  to the surface at the point  $p$ , with a sign given by the orientation of  $M$  at  $p$ .

In order to interpret the quadratic form  $\Pi_p$ , defined in (2.20), let us consider a curve parametrized by arc length  $\gamma: (-\epsilon, \epsilon) \rightarrow M$ , with  $\gamma(0) = p$ . Denoting by  $N(s) = N(\gamma(s))$ , one has  $\langle N(s), \gamma'(s) \rangle = 0$ . Thus,

$$\langle N(s), \gamma''(s) \rangle = -\langle N'(s), \gamma'(s) \rangle.$$

Therefore,

$$\begin{aligned} \Pi_p(v) &= \langle A\gamma'(0), \gamma'(0) \rangle = -\langle dN(p) \cdot \gamma'(0), \gamma'(0) \rangle \\ &= -\langle N'(0), \gamma'(0) \rangle = \langle N(0), \gamma''(0) \rangle \\ &= \langle N, kn \rangle(p) = k_n(p). \end{aligned} \tag{2.22}$$

That is, the value of the quadratic form  $\Pi_p$  in a unit vector  $v \in T_p M$  is equal to the normal curvature of a curve parametrized by arc length through  $p$  and tangent to  $v$ . In particular,

**Corollary 2.4.1** (Meusnier). All curves lying on a regular surface  $M$  and having at a given point  $p \in M$  the same tangent line have at this point the same normal curvatures.

Given an orientable regular surface  $M$  in  $\mathbb{R}^3$  and fixed a point  $p \in M$ , consider the shape operator  $A = -dN(p)$  of  $M$  at  $p$  and the corresponding

orthonormal basis  $\{e_1, e_2\}$  of  $T_p M$  that diagonalizes  $A$  as in (2.19). The numbers  $k_1$  and  $k_2$  ( $k_1 \geq k_2$ ) are the maximum and minimum, respectively, of the quadratic form  $\Pi_p$ , restricted to the unit circle of  $T_p M$ . In other words, according to (2.22),  $k_1$  and  $k_2$  are the extreme values of the normal curvatures at  $p$ . Such extreme values  $k_1$  and  $k_2$  are called the *principal curvatures* of  $M$  at  $p$ ; the corresponding eigenvectors  $e_1, e_2$  are called the *principal directions* at  $p$ .

The knowledge of the principal curvatures of  $M$  at  $p$  allows us to compute the normal curvature along any given unit vector  $v \in T_p M$ . In fact, write

$$v = \cos \theta \cdot e_1 + \sin \theta \cdot e_2,$$

where  $\theta$  is the angle between  $e_1$  and  $v$  in the orientation of  $T_p M$ . Thus, it follows from (2.22) that the normal curvature at  $p$  along  $v$  is given by

$$\begin{aligned} k_n(p) &= \Pi_p(v) = -\langle dN(p) \cdot v, v \rangle = \langle Av, v \rangle \\ &= \langle k_1 \cos \theta \cdot e_1 + k_2 \sin \theta \cdot e_2, \cos \theta \cdot e_1 + \sin \theta \cdot e_2 \rangle \\ &= k_1 \cos^2 \theta + k_2 \sin^2 \theta. \end{aligned} \quad (2.23)$$

The expression (2.23) is known as the *Euler formula*.

This description, as well as the one that follows, was introduced by Euler. Given an orientable regular surface  $M$  and a point  $p \in M$ , fix a unit vector  $v \in T_p M$ . The plane  $\pi_v$ , determined by  $v$  and the unit vector  $N(p)$  normal to  $M$ , intersects the surface  $M$  along a plane curve  $\gamma_v$ , called the *normal section* of  $M$  at  $p$  along  $v$ . Since the normal vector  $n$  is orthogonal to  $\gamma'_v(0)$  and is contained in the plane  $\pi_v$ , it follows that  $n$  and  $N$  are parallel. Therefore, the curvature  $k_v$  of  $\gamma_v$  at  $p$  is equal to the absolute value of the normal curvature at  $p$  along  $v$ . Thus, the principal curvatures  $k_1$  and  $k_2$  are the maximum and minimum values of the curvatures  $k_v$ , with  $\|v\| = 1$ .

From Examples 2.3.2 and 2.3.3, the principal curvatures of a plane  $\mathcal{P}$  and the unit sphere  $\mathbb{S}^2$  are constant and equal to 0 and 1, respectively. From Example 2.3.4, it follows that the principal curvatures of the cylinder  $\mathcal{C}$  are equal to 0 and 1 at all points. Finally, from Example 2.3.5, the principal curvatures of the hyperbolic paraboloid  $\mathcal{H}$  at the point  $(0, 0, 0)$  are equal to  $-2$  and  $2$ .

A point  $p \in M$  is called an *umbilical point* of the surface  $M$  if the principal curvatures  $k_1$  and  $k_2$  are equal at  $p$ . For instance, all the points of a sphere and a plane are umbilical points. We will see in the next section that the only surfaces which contain only umbilical points are spheres and planes.

**Definition 2.4.2.** A regular connected curve  $\gamma: I \rightarrow M$  is said to be a *line of curvature* of  $M$  if all the tangent lines of  $\gamma$  are principal direction of  $M$ .

According to Example 2.3.4, in the cylinder  $\mathcal{C}$  the generators and the equators are lines of curvature of  $\mathcal{C}$ .

**Proposition 2.4.3.** A regular connected curve  $\gamma: I \rightarrow M$ , parametrized by  $\gamma(t)$ , is a line of curvature of  $M$  if and only if

$$N'(t) = a(t)\gamma'(t),$$

where  $N(t) = N(\gamma(t))$  and  $a: I \rightarrow \mathbb{R}$  is a differentiable curve.

*Proof.* If  $\gamma'(t)$  is contained in a principal direction, then  $\gamma'(t)$  is an eigenvector of  $dN$  and

$$dN(\gamma'(t)) = N'(t) = a(t)\gamma'(t).$$

The converse is immediate. □

### Exercises

1. Show that if a regular surface is tangent to a plane along a curve, then at the points of this curve, at most one of the principal curvatures is non-zero.
2. Show that the sum of the normal curvatures for any pair of orthogonal directions, at a point  $p \in M$ , is constant.
3. Describe the region of the unit sphere covered by the image of the Gauss map of the paraboloid of revolution  $z = x^2 + y^2$ .
4. Show that the meridians of a torus are lines of curvature.
5. Determine the lines of curvature of the surface given by  $z = xy$ .
6. Let  $M_1$  and  $M_2$  be regular surfaces that intersect along a regular curve  $\gamma$  and make an angle  $\theta(p)$ , with  $p \in \gamma$ . Assume that  $\gamma$  is a line of curvature of  $M_1$ . Prove that  $\theta(p)$  is constant if and only if  $\gamma$  is a line of curvature of  $M_2$ .

## 2.5 The curvature of a regular surface

Let  $M$  be a orientable regular surface in  $\mathbb{R}^3$ . As seen in Section 2.3, for which point  $p \in M$ , the differential  $A = -dN(p)$  of the Gauss map  $N: M \rightarrow \mathbb{S}^2$  is a self-adjoint linear operator. Consider the orthonormal basis  $\{e_1, e_2\} \subset T_p M$  as in (2.19).

**Definition 2.5.1** (Euler). The *Gaussian curvature*  $K(p)$  of  $M$  at  $p$  and the *mean curvature* of  $M$  at  $p$  are defined by

$$K(p) = k_1 \cdot k_2 \quad \text{and} \quad H(p) = \frac{k_1 + k_2}{2},$$

where  $k_1$  and  $k_2$  are the principal curvatures of  $M$  at  $p$ .

In other words, the Gaussian curvature of  $M$  at  $p$  is the determinant of the shape operator  $A = -dN(p)$ , while the mean curvature of  $M$  at  $p$  is the half of the trace of  $A$ . Note that if we change the orientation of  $M$ , the determinant does not change sign; the fact that the dimension is even is essential here. However, the trace changes sign.

Let us look at some examples.

**Example 2.5.2.** At a plane  $\mathcal{P}$  in  $\mathbb{R}^3$ , the Gaussian curvature is zero, as its principal curvatures vanish identically. This is also the case for the cylinder, because its principal curvatures satisfy  $k_1 = 1$  and  $k_2 = 0$ . The unit sphere  $\mathbb{S}^2$ , however, has Gaussian curvature  $K \equiv 1$ , since its principal curvatures satisfy  $k_1 = k_2 = 1$ .

**Example 2.5.3.** Consider a regular surface  $M$  which is the graph of a differentiable function  $f: U \rightarrow \mathbb{R}$ , defined on an open set  $U \subset \mathbb{R}^2$ . The map  $\varphi: U \rightarrow \mathbb{R}^3$ , given by

$$\varphi(u, v) = (u, v, f(u, v)),$$

is a parametrization of  $M = \text{Gr}(f)$ , and thus

$$N(u, v) = \frac{\varphi_u \times \varphi_v}{\|\varphi_u \times \varphi_v\|} = \frac{1}{\sqrt{1 + f_u^2 + f_v^2}}(-f_u, -f_v, 1)$$

is a unit vector field, normal to  $M$ . In order to calculate the Gaussian curvature of  $M$ , we need to calculate the entries of the matrix of the shape operator  $A = -dN(p)$ , where  $p = \varphi(u, v)$ . We have

$$\langle A\varphi_u, \varphi_u \rangle = -\langle dN(p) \cdot \varphi_u, \varphi_u \rangle = -\langle N_u, \varphi_u \rangle = \langle N, \varphi_{uu} \rangle.$$

Since  $\varphi_{uu} = (0, 0, f_{uu})$ , we obtain

$$\langle A\varphi_u, \varphi_u \rangle = \frac{1}{\sqrt{1 + f_u^2 + f_v^2}} f_{uu}.$$

Analogously, we have

$$\langle A\varphi_v, \varphi_v \rangle = \langle N, \varphi_{vv} \rangle = \frac{1}{\sqrt{1 + f_u^2 + f_v^2}} f_{vv}$$

and

$$\langle A\varphi_u, \varphi_v \rangle = \langle N, \varphi_{uv} \rangle = \frac{1}{\sqrt{1 + f_u^2 + f_v^2}} f_{uv}.$$

Therefore,

$$A = \frac{1}{\sqrt{1 + f_u^2 + f_v^2}} \begin{pmatrix} f_{uu} & f_{uv} \\ f_{vu} & f_{vv} \end{pmatrix} = \frac{1}{\sqrt{1 + f_u^2 + f_v^2}} \text{Hess}(f).$$

It follows, in particular, that the Gaussian curvature of  $M$  is given by

$$K(p) = \det(A) = \frac{f_{uu}f_{vv} - f_{uv}^2}{(1 + f_u^2 + f_v^2)^2}.$$

## 2.6 The curvature in local coordinates

In this section we will obtain the expressions of the Gaussian curvature and of the mean curvature in a coordinate system. All parametrizations  $\varphi: U \rightarrow \varphi(U)$  considered here are assumed to be compatible with the orientation  $N$  of  $M$ . That is, in the coordinate neighborhood  $\varphi(U)$ , one has

$$N = \frac{\varphi_u \times \varphi_v}{\|\varphi_u \times \varphi_v\|}.$$

Given a point  $p \in M$ , consider a parametrization  $\varphi: U \rightarrow \varphi(U)$  of  $M$ , with  $p = \varphi(u, v)$ . Recall that the set  $\{\varphi_u, \varphi_v\}$  is a basis of  $T_p M$ . Consider a parametrized curve

$$\gamma(t) = \varphi(u(t), v(t))$$

on  $M$ , with  $\gamma(0) = p$ . The tangent vector to  $\gamma$  at  $p$  is given by

$$\gamma'(t) = \varphi_u \cdot u'(t) + \varphi_v \cdot v'(t),$$

and

$$dN(p) \cdot \gamma'(t) = N_u \cdot u'(t) + N_v \cdot v'(t).$$

Since  $N_u$  and  $N_v$  belong to  $T_p M$ , we may write

$$\begin{aligned} N_u &= a_{11}\varphi_u + a_{21}\varphi_v \\ N_v &= a_{12}\varphi_u + a_{22}\varphi_v \end{aligned} \tag{2.24}$$

and therefore,

$$dN(p) \cdot \gamma'(t) = (a_{11}u' + a_{12}v')\varphi_u + (a_{21}u' + a_{22}v')\varphi_v.$$

In another way,

$$dN(p) \begin{pmatrix} u' \\ v' \end{pmatrix} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} u' \\ v' \end{pmatrix}.$$

This shows that in the basis  $\{\varphi_u, \varphi_v\}$ , the differential  $dN(p)$  is given by the matrix  $(a_{ij})$ ,  $1 \leq i, j \leq 2$ . On the other hand, the expression of the shape operator in the basis  $\{\varphi_u, \varphi_v\}$  is given by

$$\begin{aligned} \langle A\gamma', \gamma' \rangle &= -\langle dN(\gamma'), \gamma' \rangle \\ &= -\langle N_u u' + N_v v', \varphi_u u' + \varphi_v v' \rangle \\ &= e(u')^2 + 2f u' v' + g(v')^2, \end{aligned}$$

where

$$\begin{aligned} e &= -\langle N_u, \varphi_u \rangle, \\ f &= -\langle N_v, \varphi_u \rangle = -\langle N_u, \varphi_v \rangle, \\ g &= -\langle N_v, \varphi_v \rangle. \end{aligned}$$

We will now obtain the values of  $a_{ij}$  in terms of the coefficients  $e, f, g$ . From equation (2.24), we have:

$$\begin{aligned} -e &= \langle N_u, \varphi_u \rangle = a_{11}E + a_{21}F, \\ -f &= \langle N_u, \varphi_v \rangle = a_{11}F + a_{21}G, \\ -f &= \langle N_v, \varphi_u \rangle = a_{12}E + a_{22}F, \\ -g &= \langle N_v, \varphi_v \rangle = a_{12}F + a_{22}G, \end{aligned} \tag{2.25}$$

where  $E, F$  and  $G$  denote the coefficients of the first fundamental form in the basis  $\{\varphi_u, \varphi_v\}$ , given by

$$\begin{aligned} E &= \langle \varphi_u, \varphi_u \rangle, \\ F &= \langle \varphi_u, \varphi_v \rangle, \\ G &= \langle \varphi_v, \varphi_v \rangle. \end{aligned}$$

Relations (2.25) may be expressed in matrix form by

$$-\begin{pmatrix} e & f \\ f & g \end{pmatrix} = \begin{pmatrix} a_{11} & a_{21} \\ a_{12} & a_{22} \end{pmatrix} \begin{pmatrix} E & F \\ F & G \end{pmatrix}. \tag{2.26}$$

The matrix

$$\begin{pmatrix} E & F \\ F & G \end{pmatrix}$$

is invertible, and its inverse is given by

$$\begin{pmatrix} E & F \\ F & G \end{pmatrix}^{-1} = \frac{1}{EG - F^2} \begin{pmatrix} G & -F \\ -F & E \end{pmatrix} \quad (2.27)$$

Thus, using (2.27), it follows from (2.26) that

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \frac{-1}{EG - F^2} \begin{pmatrix} e & f \\ f & g \end{pmatrix} \begin{pmatrix} G & -F \\ -F & E \end{pmatrix}. \quad (2.28)$$

Taking the determinant in the equation (2.28), we obtain

$$K(p) = \det(a_{ij}) = \frac{eg - f^2}{EG - F^2}. \quad (2.29)$$

In order to compute the mean curvature  $H$ , note that the principal curvatures  $k_1$  and  $k_2$  satisfy the equation

$$dN(p) \cdot v = -kv = -kIv,$$

for some  $v \in T_pM$ , with  $v \neq 0$ , where  $I$  denotes the identity linear map. It follows that the linear map  $dN(p) + kI$  is not invertible, and thus

$$\det \begin{pmatrix} a_{11} + k & a_{12} \\ a_{21} & a_{22} + k \end{pmatrix} = 0,$$

or,

$$k^2 + k(a_{11} + a_{22}) + a_{11}a_{22} - a_{21}a_{12} = 0. \quad (2.30)$$

The coefficients  $(a_{ij})$  of the matrix of  $dN$  in the basis  $\{\varphi_u, \varphi_v\}$  are given by

$$\begin{aligned} a_{11} &= \frac{fF - eG}{EG - F^2}, \\ a_{12} &= \frac{gF - fG}{EG - F^2}, \\ a_{21} &= \frac{eF - fE}{EG - F^2}, \\ a_{22} &= \frac{fF - gE}{EG - F^2}. \end{aligned}$$

Thus, since  $k_1$  and  $k_2$  are the roots of the quadratic equation (2.30), we obtain

$$H = \frac{1}{2}(k_1 + k_2) = -\frac{1}{2}(a_{11} + a_{22}) = \frac{1}{2} \frac{eG - 2fF + gE}{EG - F^2}. \quad (2.31)$$

**Remark 2.6.1.** From (2.30) and (2.31), we conclude that

$$k^2 - 2Hk + K = 0$$

and therefore,

$$k = H \pm \sqrt{H^2 - K}. \quad (2.32)$$

From relation (2.32), it follows that if we choose  $k_1(p) \geq k_2(p)$ , with  $p \in M$ , then the functions  $k_1$  and  $k_2$  are continuous in  $M$ . In fact,  $k_1$  and  $k_2$  are differentiable in  $M$ , except perhaps at the umbilical points of  $M$ .

### Exercises

1. Consider a parametrization of a regular surface  $M$  such that  $F = f = 0$ . Show that the principal curvatures are given by  $k_1 = e/E$  and  $k_2 = g/G$ .

2. An *asymptotic direction* of a regular surface  $M$  at a point  $p \in M$  is a direction of  $T_pM$  for which the normal curvature is zero. An *asymptotic curve* of  $M$  is a regular connected curve  $\gamma \subset M$  such that for each point  $p \in \gamma$ , the tangent line of  $\gamma$  at  $p$  is an asymptotic direction.

- (a) Consider a parametrization  $\varphi$  of  $M$ . Show that a connected regular curve  $\gamma: I \rightarrow M$  in the coordinate neighborhood of  $\varphi(u, v)$  is an asymptotic curve of  $M$  if and only if, for any parametrization  $\gamma(t) = \varphi(u(t), v(t))$  of  $\gamma$ , one has

$$e(u')^2 + 2fu'v' + g(v')^2 = 0, \quad t \in I. \quad (2.33)$$

Equation (2.33) is called the *differential equation of the asymptotic curves*.

- (b) Determine the asymptotic curves of the surface given by  $z = xy$ .

3. Consider the *Enneper surface*  $M$  parametrized by

$$\varphi(u, v) = \left( u - \frac{u^3}{3} + uv^2, v - \frac{v^3}{3} + vu^2, u^2 - v^2 \right).$$

- (a) Show that the coefficients of the first e second fundamental form of  $M$  are given by

$$E = G = (1 + u^2 + v^2)^2, \quad F = 0$$

and

$$e = 2, \quad f = 0, \quad g = -2,$$

respectively.

- (b) Determine the principal curvatures of  $M$ .
- (c) Show that the asymptotic curves are  $u + v = \text{const}$  and  $u - v = \text{const}$ .

## 2.7 Rotational surfaces

In this section we will study in detail a given rotational surface  $M$ , which is parameterized by

$$\varphi(u, v) = (f(u) \cos v, f(u) \sin v, g(u)),$$

where  $a < u < b$  and  $0 < v < 2\pi$ . It follows from Exercise ?? that  $\varphi$  is in fact a parameterization for  $M$ . Let us assume that the generator curve  $\gamma = (f(u), 0, g(u))$  is parameterized by arc length, that is,

$$f'(u)^2 + g'(u)^2 = 1. \quad (2.34)$$

Then,

$$\varphi_u = (f' \cos v, f' \sin v, g') \quad \text{and} \quad \varphi_v = (-f \sin v, f \cos v, 0),$$

such that

$$E = 1, \quad F = 0, \quad G = f^2.$$

Moreover, we have

$$\begin{aligned} \varphi_{uu} &= (f'' \cos v, f'' \sin v, g'') \\ \varphi_{uv} &= (-f' \sin v, f' \cos v, 0) \\ \varphi_{vv} &= (-f \cos v, -f \sin v, 0) \end{aligned}$$

and

$$N = \frac{\varphi_u \times \varphi_v}{\|\varphi_u \times \varphi_v\|} = (-g' \cos v, -g' \sin v, f').$$

Thus,

$$\begin{aligned} l &= \langle \varphi_{uu}, N \rangle = f'g'' - f''g' \\ m &= \langle \varphi_{uv}, N \rangle = 0 \\ n &= \langle \varphi_{vv}, N \rangle = fg'. \end{aligned}$$

Therefore, the Gaussian curvature of  $M$  is given by

$$K = \frac{ln - m^2}{EG - F^2} = \frac{(f'g'' - f''g')g'}{f} = k_\gamma \frac{g'}{f}, \quad (2.35)$$

where  $k_\gamma = f'g'' - f''g'$  is the curvature of  $\gamma$ . Sometimes it is convenient to rewrite the Gaussian curvature  $K$  in another way. Differentiation of (2.34) gives  $f'f'' + g'g'' = 0$ . Substituting this identity into (2.35) yields

$$K = -\frac{f''}{f}.$$

In this case, since  $F = m = 0$ , it follows from Exercise 2.6.1 that the principal curvatures of  $M$  are  $k_1 = k_\gamma$  and  $k_2 = \frac{g'}{f}$ . Finally, the mean curvature of  $M$  is given by

$$H = \frac{1}{2} \left( \frac{g''}{f'} + \frac{g'}{f} \right) = \frac{(fg')'}{(f^2)'}.$$
 (2.36)

**Example 2.7.1.** A particular case of rotational surface is the torus parametrized by

$$\varphi(u, v) = ((a + r \cos u) \cos v, (a + r \cos u) \sin v, r \sin u),$$

with  $0 < u, v < 2\pi$  and  $0 < r < a$ . In this case, the Gaussian curvature of the torus is given by

$$K = \frac{r \cos u}{a + r \cos u}.$$
 (2.37)

From the expression (2.37), it follows that  $K = 0$  along the parallels  $u = \pi/2$  and  $u = 3\pi/2$ . The points of the torus given by  $\pi/2 < u < 3\pi/2$ ,  $K$  is negative. In the region given by  $0 < u < \pi/2$  or  $3\pi/2 < u < 2\pi$ , the Gaussian curvature is positive.

**Definition 2.7.2.** A point  $P$  of a regular surface  $M$  is called

- (1) *Elliptic* if  $\det(dN(p)) > 0$ .
- (2) *Hyperbolic* if  $\det(dN(p)) < 0$ .
- (3) *Parabolic* if  $\det(dN(p)) = 0$ , with  $dN(p) \neq 0$ .
- (4) *Planar* if  $dN(p) = 0$ .

We point out that this classification does not depend on the choice of the orientation of  $M$ . At an elliptic point the Gaussian curvature is positive. That is, both principal curvatures have the same sign, and therefore all curves passing through this point have their normal vectors pointing toward the same side of the tangent plane.

At a hyperbolic point  $p \in M$ , the Gaussian curvature is negative. The principal curvatures have opposite signs, and therefore there are curves

through  $p$  whose normal vectors at  $p$  point toward any of the sides of the tangent plane at  $p$ .

At a parabolic point, the Gaussian curvature is zero, but one of the principal curvatures is not zero. Finally, at a planar point, all principal curvatures are zero.

### Exercises

1. Consider the surface obtained by rotating the curve  $y = x^3$  around the line  $x = 1$ , with  $-1 < x < 1$ . Show that the points obtained by rotation of the origin  $(0, 0)$  of the curve are planar points of the surface.
2. Show that a surface which is compact has an elliptic point.

## 2.8 Minimal surfaces

Let  $M$  be an orientable regular surface.

**Definition 2.8.1.** We say that  $M$  is a *minimal surface* if the mean curvature  $H$  vanishes everywhere.

Historically, this concept is related to the problem of characterizing the surface with smallest area spanned by a given boundary, a problem raised by Lagrange in 1760. A simple example of minimal surface is the plane, since both principal curvatures are identically zero.

**Example 2.8.2.** Another example of minimal surface is the *catenoid*, which is generated by the rotation of the catenary  $y = a \cosh \frac{z}{a}$  around the  $z$  axis, where  $a > 0$ . Thus,

$$\varphi(u, v) = \left( a \cosh \frac{u}{a} \cos v, a \cosh \frac{u}{a} \sin v, u \right)$$

is a parametrization for the catenoid, where  $0 < v < 2\pi$  and  $u \in \mathbb{R}$ . We have

$$\begin{aligned} \varphi_u &= \left( \sinh \frac{u}{a} \cos v, \sinh \frac{u}{a} \sin v, 1 \right), \\ \varphi_v &= \left( -a \cosh \frac{u}{a} \sin v, a \cosh \frac{u}{a} \cos v, 0 \right). \end{aligned}$$

Thus,

$$E = \cosh^2 \frac{u}{a}, \quad F = 0, \quad G = a^2 \cosh^2 \frac{u}{a}.$$

Moreover,

$$\begin{aligned}\varphi_{uu} &= \left( \frac{1}{a} \cosh \frac{u}{a} \cos v, \frac{1}{a} \cosh \frac{u}{a} \sin v, 0 \right), \\ \varphi_{uv} &= \left( -\sinh \frac{u}{a} \sin v, \sinh \frac{u}{a} \cos v, 0 \right), \\ \varphi_{vv} &= \left( -a \cosh \frac{u}{a} \cos v, -a \cosh \frac{u}{a} \sin v, 0 \right),\end{aligned}$$

and

$$N(u, v) = \frac{1}{\cosh \frac{u}{a}} \left( -\cos v, -\sin v, \sinh \frac{u}{a} \right).$$

Thus, the coefficients of the second fundamental form are given by

$$e = -\frac{1}{a}, \quad f = 0, \quad g = a.$$

Therefore, the principal curvatures of the catenoid are given by

$$k_1 = -\frac{1}{a} \operatorname{sech}^2 \frac{u}{a} \quad \text{and} \quad k_2 = \frac{1}{a} \operatorname{sech}^2 \frac{u}{a},$$

which implies that the mean curvature  $H$  vanishes everywhere.




Figure 2.1: The catenoid.

**Proposition 2.8.3.** The only minimal rotational surfaces are the plane and the catenoid.

*Proof.* Let  $M \subset \mathbb{R}^3$  be a minimal rotational surface, parametrized by

$$\varphi(u, v) = (f(u) \cos v, f(u) \sin v, g(u)),$$

where  $a < u < b$ ,  $0 < v < 2\pi$ , and the generator curve  $\gamma = (f(u), 0, g(u))$  is parameterized by arc length, that is,

$$f'(u)^2 + g'(u)^2 = 1. \tag{2.38}$$

The mean curvature  $H$  of  $M$  is

$$H = \frac{1}{2} \left( k_\gamma + \frac{g'}{f} \right), \quad (2.39)$$

where  $k_\gamma = f'g'' - f''g'$  is the curvature of  $\gamma$ . If  $f' = 0$ , it follows from (2.39) and the fact that  $H = 0$  that  $g' = 0$ , which is a contradiction with (2.38). Therefore we can assume that  $f'$  is never zero. On the other hand, it follows from (2.36) that  $(fg')' = 0$ . That is, we need to solve the equation

$$fg' = k, \quad (2.40)$$

where  $k$  is a constant. From (2.38), we have

$$g' = \pm \sqrt{1 - (f')^2}. \quad (2.41)$$

Replacing (2.41) into (2.40), one has

$$f' = \pm \sqrt{1 - (k/f)^2}. \quad (2.42)$$

Equation (2.42) can be rewritten as

$$\frac{f}{\sqrt{f^2 - k^2}} df = \pm du,$$

and integrating we obtain

$$f(u) = \pm \sqrt{k^2 + (u + c)^2},$$

for some constant  $c$ . This constant  $c$  can be chosen to be zero by redefining the instant  $u = 0$ . Since  $f > 0$ , we obtain

$$f(u) = \sqrt{u^2 + k^2}.$$

If  $k = 0$  then  $f(u) = \pm u$ . Thus  $g(u) = d$  is constant, and the surface can be parametrized as

$$\varphi(u, v) = (u \cos v, u \sin v, d),$$

which corresponds to the case of the plane. Suppose now  $k \neq 0$ . By integrating (2.40) we

$$g(u) = k \ln(u + \sqrt{u^2 + k^2}) + c.$$

We choose the constant  $c$  such that  $g(0) = 0$ , that is,  $c = -k \ln |k|$ . We may assume that  $k > 0$ . In fact, changing the sign of  $k$  is equivalent to changing

the sign of  $g$ , which corresponds to a reflection on the plane  $z = 0$ . Thus, we can write  $g$  as

$$g(u) = k \ln(u/k + \sqrt{(u/k)^2 + 1}).$$

Finally, making the change of variable  $t = g(u)$ , we get

$$\gamma(t) = (k \cosh \frac{t}{k}, 0, t),$$

which is a catenary. □

**Example 2.8.4.** Consider a helix in  $\mathbb{R}^3$  given by

$$\gamma(u) = (\cos u, \sin u, au), \quad a > 0.$$

Through each point of the helix, draw a line parallel to the  $xy$  plane and intersecting the  $z$  axis. The surface generated by these lines is called a *helicoid* and can be parametrized by

$$\varphi(u, v) = (v \cos u, v \sin u, au),$$

with  $0 < u < 2\pi$  and  $v \in \mathbb{R}$ . We have

$$\varphi_u = (-v \sin u, v \cos u, a) \quad \text{and} \quad \varphi_v = (\cos u, \sin u, 0).$$

Thus, the coefficients of the first fundamental form are given by

$$E = a^2 + v^2, \quad F = 0, \quad G = 1,$$

and

$$N = \frac{\varphi_u \times \varphi_v}{\|\varphi_u \times \varphi_v\|} = \frac{1}{\sqrt{a^2 + v^2}}(-a \sin u, a \cos u, -v)$$

is a unit normal vector field to the helicoid. Moreover,

$$\begin{aligned} \varphi_{uu} &= (-v \cos u, -v \sin u, 0), \\ \varphi_{uv} &= (-\sin u \cos u, 0), \\ \varphi_{vv} &= (0, 0, 0), \end{aligned}$$

which imply that the coefficients of the second fundamental form are given by

$$e = 0, \quad f = \frac{a}{\sqrt{a^2 + v^2}}, \quad g = 0,$$

Therefore, we conclude that  $H = 0$ .




Figure 2.2: The helicoid.

### Exercises

1. Determine the asymptotic curves and the lines of curvature of the helicoid parametrized by

$$\varphi(u, v) = (v \cos u, v \sin u, au),$$

and of the catenoid parametrized by

$$\varphi(u, v) = (\cosh u \cos v, \cosh u \sin v, u).$$

2. Prove that there are no compact minimal surfaces.

## 2.9 Normal variations

Let  $M \subset \mathbb{R}^3$  be an orientable regular surface and let  $f: M \rightarrow \mathbb{R}$  be a smooth function.

**Definition 2.9.1.** A *normal variation* of  $M$  determined by  $f$  is a one-parameter family of surfaces  $M^t$ , with  $t \in (-\epsilon, \epsilon)$ , given by

$$p^t = p + tf(p)N(p),$$

where  $N$  is the Gauss map of the surface  $M$ .

Para  $\epsilon > 0$  sufficiently small, each set  $M^t$  is a regular surface, called a *variation of the surface*. In fact, if  $\varphi$  is a parametrization of  $M$ , then

$$\varphi^t = \varphi + tfN$$

is parametrization of  $M^t$ , with  $t \in (-\epsilon, \epsilon)$ . Note that  $M^0 = M$ , and when  $f \equiv 1$ ,  $M^t$  is a *parallel surface* to  $M$  at a distance  $t$ .

Let  $M^t$  be a normal variation of  $M$ , with  $t \in (-\epsilon, \epsilon)$ , determined by a smooth function  $f: M \rightarrow \mathbb{R}$ . Given a compact domain  $D \subset M$ , set

$$D^t = \{p^t \in M^t : p \in D\}$$

the corresponding domain in  $M^t$ . For each instant  $t \in (-\epsilon, \epsilon)$ , define

$$A(t) = \text{Area}(D^t).$$

**Theorem 2.9.2.** *For any normal variation of  $M$  determined by a smooth function  $f: M \rightarrow \mathbb{R}$ , and for any compact domain  $D \subset M$ , one has*

$$A'(0) = -2 \iint_D H f dA. \quad (2.43)$$

The concept of minimal surface is related to the following problem proposed by Lagrange in 1760: *Given a smooth closed curve  $\gamma$ , determine the surface of minimum area that has this curve as its boundary.*




Figure 2.3: A soap bubble from the curve.

Suppose there is a surface  $M$  that there exists a solution to Lagrange problem. Consider a normal variation  $M^t$  for  $M$ , with  $t \in (-\epsilon, \epsilon)$ , determined by a smooth function  $f: M \rightarrow \mathbb{R}$ , with  $f|_\gamma = 0$ . This means that the variation leaves the curve  $\gamma$  fixed. Since the area of  $M$  is minimal we obtain, in particular, that

$$A(t) \geq A(0),$$

for every  $t \in (-\epsilon, \epsilon)$  and for any such variation. Therefore, we have  $A'(0) = 0$ , for any smooth function  $f: M \rightarrow \mathbb{R}$ , with  $f|_\gamma = 0$ .

**Proposition 2.9.3.** A surface  $M$  is minimal if and only if  $A'(0) = 0$ , for every such normal variation of  $M$ , and every compact domain  $D \subset M$ .

*Proof.* If  $H \equiv 0$ , it follows directly from (2.43) that  $A'(0) = 0$ . Conversely, suppose that  $A'(0) = 0$  for any normal variation determined by a smooth function  $f: M \rightarrow \mathbb{R}$ , with  $f|_\gamma = 0$ , and suppose that there exists a point

$p \in M$  such that  $H(p) > 0$ . Choose such  $f$  such that  $f \geq 0$ ,  $f(p) = H(p)$ , and  $f|_{M \setminus D} = 0$ , where  $D \subset M$  is a compact domain around  $p$ , with  $H > 0$ . For such a function  $f$ , we have

$$A'(0) = -2 \iint_D H f dA < 0,$$

a contradiction. Therefore, one has  $H \equiv 0$ .  $\square$

It follows from Proposition 2.9.3 that, if there exists a surface  $M$  of minimum area, then  $H \equiv 0$ . That is, surfaces of minimum area are minimal surfaces in the sense of Definition 2.8.1. The converse, however, is false.

**Example 2.9.4.** Consider a bubble soap that is in equilibrium under the action of the surface tension of the liquid. We claim that this equilibrium surface has mean curvature  $H \equiv 0$ . In fact, it follows from a formula, due to Laplace, establishing that the pressure at each point carried by the surface on the environment is carried out along the normal to the surface and proportional to  $H$ . Since the surface is in equilibrium, such pressure zero, and therefore  $H \equiv 0$ .




Figure 2.4: A soap bubble.

**Example 2.9.5.** Consider a surface  $M$  in  $\mathbb{R}^3$  which is the graph of a smooth function  $z = f(x, y)$ . The condition  $H \equiv 0$  is equivalent to the following equation:

$$(1 + f_y^2)f_{xx} + 2f_x f_y f_{xy} + (1 + f_x^2)f_{yy} = 0. \quad (2.44)$$

For example, the linear functions

$$f(x, y) = ax + by + c,$$

where  $a, b, c \in \mathbb{R}$ , are trivial solutions of (2.44).

**Example 2.9.6** (Scherk 1835). Suppose, in the equation (2.44), that  $f$  can be written as

$$f(x, y) = g(x) + h(y).$$

In this case, equation (2.44) becomes

$$(1 + h'(y)^2)g''(x) + (1 + g'(x)^2)h''(y) = 0,$$

that is,

$$\frac{g''(x)}{1 + g'(x)^2} = -\frac{h''(y)}{1 + h'(y)^2} = \text{constant}.$$

By integration, we obtain

$$g(x) = \ln \cos x \quad \text{and} \quad h(y) = \ln \cos y,$$

up to constants. Therefore, up to translations and dilations, a piece of the surface can be represented as the graph of the function

$$f(x, y) = \ln \frac{\cos x}{\cos y},$$

with  $0 < x, y < \frac{\pi}{2}$ .




Figure 2.5: The Scherk surface.

## Exercises

1. Prove that the Enneper surface is minimal.

## 2.10 Appendix: Regular curves

A parametrized differentiable curve  $\lambda: I \rightarrow \mathbb{R}^3$  is said to be *regular*  $\lambda'(t) \neq 0$  for all  $t \in I$ . Fixed an instant  $t_0 \in I$ , the *arc length* of  $\lambda$ , from the instant  $t_0$ , is the function

$$s(t) = \int_{t_0}^t \|\lambda'(t)\| dt. \quad (2.45)$$

We will say that  $\lambda$  is *parameterized by arc length* if for every  $t_1, t_2 \in I$ , with  $t_1 \leq t_2$ , the arc length of  $\lambda$  from  $t_1$  to  $t_2$  is equal to  $t_2 - t_1$ , that is

$$\int_{t_1}^{t_2} \|\lambda'(t)\| dt = t_2 - t_1. \quad (2.46)$$

Note that, if  $\|\lambda'(t)\| = 1$  for any  $t \in I$ , it follows directly from (2.46) that  $\lambda$  is parameterized by arc length. The converse also holds (cf. Exercise 2.4.1). For instance, the helix

$$\lambda(t) = \left( \cos \frac{t}{\sqrt{2}}, \sin \frac{t}{\sqrt{2}}, \frac{t}{\sqrt{2}} \right)$$

is a curve parametrized by arc length, since  $\|\lambda'(t)\| = 1$  for every  $t \in I$ .

**Proposition 2.10.1.** Let  $\lambda: I \rightarrow \mathbb{R}^3$  be a regular curve. Then, the arc length function (2.45) admits an inverse  $h: J \rightarrow I$ , defined on the open interval  $J = s(I)$ , and  $\beta = \lambda \circ h$  is a reparametrization of  $\lambda$  that is parameterized by arc length.

*Proof.* Since  $\lambda$  is a regular curve, it follows that the arc length function  $s$  is strictly increasing, because

$$s'(t) = \|\lambda'(t)\| = 1,$$

for every  $t \in I$ . Thus,  $s$  admits an inverse  $h: J \rightarrow I$ . Since  $h \circ s = id$ , one has  $\frac{dh}{ds} \cdot \frac{ds}{dt} = 1$ , and therefore

$$\frac{dh}{ds} = \frac{1}{s'(t)} = \frac{1}{\|\lambda'(t)\|} > 0.$$

This implies that  $\beta(s) = \lambda(h(s))$ ,  $s \in J$ , is a reparametrization of  $\lambda$  that satisfies

$$\|\beta'(s)\| = \left\| \frac{d\beta}{ds} \right\| = \left\| \frac{d\lambda}{dt} \cdot \frac{dh}{ds} \right\| = \frac{\|\lambda'(t)\|}{\|\lambda'(t)\|} = 1,$$

for every  $s \in J$ . Therefore, it follows from Exercise 2.4.1 that  $\beta$  is parameterized by arc length.  $\square$

The curve  $\beta$  given by Proposition 2.10.1 is called a *reparametrization of  $\lambda$  by arc length*. Note that this reparametrization is not uniquely determined, because it depends of the arc length function, which depends of the instant  $t_0$  fixed.

**Example 2.10.2.** Consider the helix parametrized by

$$\lambda(t) = (\cos t, \sin t, t), \quad t \in \mathbb{R}.$$

Since  $\|\lambda'(t)\| = \sqrt{2}$ , the arc length function of  $\lambda$ , from  $t_0 = 0$ , is  $s(t) = \sqrt{2}t$ , and its inverse is given by  $h(s) = \frac{1}{\sqrt{2}}s$ . Thus,

$$\beta(s) = \lambda(h(s)) = \left( \cos \frac{s}{\sqrt{2}}, \sin \frac{s}{\sqrt{2}}, \frac{s}{\sqrt{2}} \right)$$

is a reparametrization of  $\lambda$  by arc length.

Let  $\lambda: I \rightarrow \mathbb{R}^3$  be a curve parametrized by arc length  $s$ . Since  $\|\lambda'(s)\| = 1$  for every  $s \in I$ , the norm  $\|\lambda''(s)\|$  of the second derivative measures the rate of change of the tangent line to  $\lambda$ , that is, how rapidly the curve pulls away from the tangent line at  $s$ .

**Definition 2.10.3.** Let  $\lambda: I \rightarrow \mathbb{R}^3$  be a curve parametrized by arc length  $s$ . The number  $k(s) = \|\lambda''(s)\|$  is called the *curvature* of  $\lambda$  at  $s$ .

If  $\lambda$  is a straight line, parametrized by  $\lambda(s) = as + b$ , with  $\|a\| = 1$ , then  $k(s) = 0$ . Conversely, if  $k(s) = 0$  for every  $s$ , then by integration one has  $\lambda(s) = as + b$ , and the curve is a straight line. Therefore, a curve parametrized by arc length has vanishing curvature if and only if its trace is contained in a straight line.

**Example 2.10.4.** Consider a helix parametrized by

$$\lambda(s) = \left( a \cos \frac{s}{c}, a \sin \frac{s}{c}, \frac{bs}{c} \right), \quad s \in \mathbb{R},$$

where  $a, b, c$  are positive constants, with  $a^2 + b^2 = c^2$ . Note that the parameter  $s$  is the arc length, and a straightforward computation shows that  $k(s) = \frac{a}{c^2}$ , for every  $s \in \mathbb{R}$ .

At points where  $k(s) \neq 0$ , a unit vector  $n(s)$  is well defined by the equation

$$\lambda''(s) = k(s) \cdot n(s).$$

Since  $\lambda''(s)$  is orthogonal to  $\lambda'(s)$ , it follows that  $n(s)$  is orthogonal to  $\lambda'(s)$ , and it is called the *normal vector* of  $\lambda$  at  $s$ .

## Exercises

1. Show that a regular curve  $\lambda: I \rightarrow \mathbb{R}^3$  is parameterized by arc length if and only if  $\|\lambda'(t)\| = 1$  for every  $t \in I$ .
2. Assume that all normals of a parametrized curve cross through a fixed point. Prove that the trace of the curve is contained in a circle.
3. Let  $\lambda: I \rightarrow \mathbb{R}^3$  be a regular curve such that all its tangent lines cross through a fixed point. Prove that the trace of  $\lambda$  is contained in a straight line.
4. Calculate the curvature of the *catenary*, which is the curve  $\lambda: \mathbb{R} \rightarrow \mathbb{R}^2$  parametrized by  $\lambda(t) = (t, \cosh t)$ .

# Bibliography

- [1] M. P. do Carmo, *Differential Geometry of Curves and Surfaces*, Prentice-Hall, 1976.
- [2] M. Spivak, *A Comprehensive Introduction to Differential Geometry*, Vol. 3, 3rd Edition, Publish or Perish, 1999.